arXiv:1102.2047v3 [math.QA] 22 Sep 2011

A SEMINORMAL FORM FOR PARTITION ALGEBRAS

JOHN ENYANG

Contents

1. Introduction 1
2. Preliminaries 2
3. A Murphy Basis 7
4. A Seminormal Form 17
5. Main Results 20
6. Central Element Recursions 35
7. Tables of Representing Matrices 41

References 46

Abstract. Using a new presentation for partition algebras [arXiv:1009.1939], we give explicit combinatorial formulae for the seminormal representations of the partition algebras. Our results generalise to the partition algebras the classical formulae given by Young for the symmetric group.

1. Introduction

The partition algebras $A_k(n)$, for $k, n \in \mathbb{Z}_{\geqslant 0}$, are a family of algebras defined in the work of Martin and Jones in [Mar], [Mar1], [Jo] in connection with the Potts model and higher dimensional statistical mechanics. By [Jo], the partition algebra $A_k(n)$ is in Schur–Weyl duality with the symmetric group $\mathfrak{S}_n$ acting diagonally on the k–fold tensor product $V^{\otimes k}$ of its n–dimensional permutation representation V. In [Mar2], Martin defined the partition algebras $A_{k+\frac{1}{2}}(n)$ as the centralisers of the subgroup $\mathfrak{S}_{n-1} \subseteq \mathfrak{S}_n$ acting on $V^{\otimes k}$. Including the algebras $A_{k+\frac{1}{2}}(n)$ in the tower

$$A_0(n) \subseteq A_{\frac{1}{2}}(n) \subseteq A_1(n) \subseteq A_{1+\frac{1}{2}}(n) \subseteq \cdots \tag{1.1}$$

allowed for the simultaneous analysis of the whole tower of algebras (1.1) using the Jones Basic construction by Martin [Mar2] and Halverson and Ram [HR]. The partition algebras (1.1) have connections to Deligne's category $\underline{\mathrm{Rep}}(S_t)$ [CO], and are important examples of cellular algebras [Xi],[DW],[GG].

Halverson and Ram [HR] used Schur–Weyl duality to show that certain diagrammatically defined elements in the partition algebras play an analogous role to the classical Jucys–Murphy elements in the symmetric group. By definition, a seminormal form for $A_k(n)$ or $A_{k+\frac{1}{2}}(n)$is an irreducible matrix representation relative to a basis of eigenvectors for Jucys–Murphy elements.

Classically, the seminormal form appeared in Young's construction [Yo] of irreducible representations for the symmetric group $\mathfrak{S}_k$ (see [Kl] or [VO] for a modern treatment of the subject). Kosuda [Ko1] has used the presentation in [HR] to compute the seminormal representations for the partition algebra $A_3(n)$. The seminormal representations of the subalgebra of the partition

---

Key words and phrases. Partition algebra; cellular algebra; Jucys–Murphy element; seminormal form.

algebra $A_k(n)$ that acts as centraliser of $G(r,1,n)$ on the tensor space $V^{\otimes k}$, for $n \geqslant k$ and $r > k$, have been constructed by Kosuda [Ko].

In this paper we provide explicit combinatorial formulae for the seminormal representations of the partition algebras $A_k(n)$ and $A_{k+\frac{1}{2}}(n)$. The new approach here is to use the presentation for partition algebras in [En] to compute seminormal representations. For the representations of $A_k(n)$ and $A_{k+\frac{1}{2}}(n)$ which factor through $\mathfrak{S}_k$, our formulae coincide with those given by Young [Yo]. Our construction of seminormal representations provides a partition algebra analogue of the work of Nazarov [Na] for the Brauer algebras and the work of Leduc and Ram [LR] for Brauer and BMW algebras.

Nazarov introduced a remarkable recursion for special central elements in the Brauer algebras and established the relation between these central elements and seminormal representations (see Corollary 3.10 and Proposition 4.2 of [Na]). A similar relation was established by Beliakova and Blanchet for the BMW algebras (see Lemma 7.2 and Lemma 7.4 of [BB]). In this paper, we obtain analogous recursions for central elements in the partition algebras and explain the relation between these central elements and the seminormal representations of the partition algebras.

In §2 we recall the presentations of the partition algebras from [HR] and [En], and state the definition of the Jucys–Murphy elements from [En]. In §3 we show that the Jucys–Murphy elements act triangularly on the partition algebras relative to the Murphy–type bases given in [En1]. In §4 we define seminormal bases for the partition algebras and in §5 we state and prove explicit combinatorial formulae for the images of the generators in the seminormal matrix representations of the partition algebras. In §6 we define central elements (6.3) by contracting powers of the Jucys–Murphy elements in the partition algebras and derive partition algebra analogues of the Nazarov recursions (see Proposition 6.1). In Proposition 6.2 we establish the relation between seminormal matrix entries and the Nazarov–type recursions for central elements in the partition algebras. In §7 we give tables of representing matrices of small rank.

Acknowledgements. The author would like to express his gratitude to Arun Ram for numerous stimulating conversations throughout the course of this work. The author is also much indebted to Fred Goodman for several helpful discussions related to this research. This research was supported by the Australian Research Council (grant ARC DP–0986774) at the University of Melbourne.

2. PRELIMINARIES

2.1. **Combinatorics.** We recall the notation established in [En1]. Let k denote a non–negative integer and $\mathfrak{S}_k$ be the symmetric group acting on $\{1,\dots,k\}$ on the right. For i an integer, $1 \leqslant i < k$, let s_i denote the transposition $(i,i+1)$. Then $\mathfrak{S}_k$ is presented as a Coxeter group by generators $s_1, s_2, \dots, s_{k-1}$, with the relations

$$\begin{aligned} &s_i^2 = 1, && \text{for } i = 1,\dots,k-1,\\ &s_i s_j = s_j s_i, && \text{for } j \neq i+1.\\ &s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, && \text{for } i = 1,\dots,k-2. \end{aligned}$$

An product $w = s_{i_1} s_{i_2} \cdots s_{i_j}$ in which j is minimal is called a *reduced expression* for w and $j = \ell(w)$ is the *length* of w. If $i, j = 1,\dots,k$, define

$$w_{i,j} = \begin{cases} s_i s_{i+1} \cdots s_{j-1}, & \text{if } j \geqslant i,\\ s_{i-1} s_{i-2} \cdots s_j, & \text{if } i > j. \end{cases}$$

If $k > 0$, a *partition* of k is a non–increasing sequence $\lambda = (\lambda_1, \lambda_2, \dots)$ of integers, $\lambda_i \geqslant 0$, such that $\sum_{i\geqslant 1} \lambda_i = k$; otherwise, if $k = 0$, write $\lambda = \emptyset$ for the empty partition. The fact that λ is a partition of k will be denoted by $\lambda \vdash k$. If λ is a partition, we will also write $|\lambda| = \sum_{i\geqslant 1} \lambda_i$. The integers $\{\lambda_i \mid \text{for } i \geqslant 1\}$ are the *parts* of λ. If $\lambda \vdash k$, the *Young diagram* of λ is the set

$$[\lambda] = \{(i,j) \,|\, \lambda_i \geqslant j \geqslant 1 \text{ and } i \geqslant 1\,\} \subseteq \mathbb{N} \times \mathbb{N}.$$

The elements of $[\lambda]$ are the *nodes* of λ and more generally a node is a pair $(i,j) \in \mathbb{Z} \times \mathbb{Z}$. The diagram $[\lambda]$ is traditionally represented as an array of boxes with λ_i boxes on the i–th row. For example, if $\lambda = (3,2)$, then $[\lambda] = \begin{array}{|c|c|c|}\hline & & \\ \hline & \\ \cline{1-2}\end{array}$. We will usually identify the partition λ with its diagram and write λ in place of $[\lambda]$. Let λ be a partition. A node (i,j) is an *addable* node of λ if $(i,j) \notin \lambda$ and $\mu = \lambda \cup \{(i,j)\}$ is a partition; in this case (i,j) is also referred to as a *removable* node of μ. Let $A(\lambda)$ and $R(\lambda)$ respectively denote the set of addable nodes and removable nodes of λ.

The dominance $\trianglerighteq$ on partitions of k is defined as follows: if $\lambda \vdash k$ and $\mu \vdash k$, then $\lambda \trianglerighteq \mu$ if

$$\textstyle\sum_{i=1}^{j} \lambda_i \geqslant \sum_{i=1}^{j} \mu_i \qquad\qquad \text{for all } j \geqslant 1.$$

We write $\lambda \triangleright \mu$ to mean that $\lambda \trianglerighteq \mu$ and $\lambda \neq \mu$.

Let $\lambda \vdash k$. A λ–tableau $\mathfrak{t}$ from the nodes of the diagram $[\lambda]$ to the integers $\{1, 2, \dots, k\}$. A given λ–tableau $\mathfrak{t} : [\lambda] \to \{1, 2, \dots, k\}$ can be represented by labelling the nodes of the diagram $[\lambda]$ with the integers $1, 2, \dots, k$. For example, if $k = 6$ and $\lambda = (3,2,1)$,

$$\mathfrak{t} = \begin{array}{|c|c|c|}\hline 1 & 4 & 6 \\ \hline 2 & 3 \\ \cline{1-2} 5 \\ \cline{1-1}\end{array} \tag{2.1}$$

represents a λ–tableau. If $\lambda \vdash k$, let $\mathfrak{t}^\lambda$ denote the λ–tableau in which $1, 2, \dots, k$ are entered in increasing order from left to right along the rows of $[\lambda]$. Thus in the previous example where $k = 6$ and $\lambda = (3,2,1)$,

$$\mathfrak{t}^\lambda = \begin{array}{|c|c|c|}\hline 1 & 2 & 3 \\ \hline 4 & 5 \\ \cline{1-2} 6 \\ \cline{1-1}\end{array}\,. \tag{2.2}$$

The tableau $\mathfrak{t}^\lambda$ is the *row reading tableau* of shape λ. The symmetric group $\mathfrak{S}_k$ acts on the set of λ–tableaux on the right by permuting the integer labels of the nodes of $[\lambda]$. For example,

$$\begin{array}{|c|c|c|}\hline 1 & 2 & 3 \\ \hline 4 & 5 \\ \cline{1-2} 6 \\ \cline{1-1}\end{array}\,(2,4)(3,6,5) \;=\; \begin{array}{|c|c|c|}\hline 1 & 4 & 6 \\ \hline 2 & 3 \\ \cline{1-2} 5 \\ \cline{1-1}\end{array}\,.$$

If $\lambda \vdash k$, the *Young subgroup* $\mathfrak{S}_\lambda$ is defined to be the row stabiliser of $\mathfrak{t}^\lambda$ in $\mathfrak{S}_k$. For instance, when $k = 6$ and $\lambda = (3,2,1)$, as in (2.2), then $\mathfrak{S}_\lambda = \langle s_1, s_2, s_4 \rangle$.

2.2. **Partition algebras.** In this section we follow the exposition given by Halverson and Ram in [HR]. For $k = 1, 2, \dots,$ let

$$\begin{aligned} A_k &= \{\text{set partitions of } \{1, 2, \dots, k, 1', 2', \dots, k'\}\}, \qquad \text{and,} \\ A_{k-\frac{1}{2}} &= \{d \in A_k \mid k \text{ and } k' \text{ are in the same block of } d\}. \end{aligned}$$

Any element $\rho \in A_k$ may be represented as a graph with k vertices in the top row, labelled from left to right, by $1, 2, \dots, k$ and k vertices in the bottom row, labelled, from left to right by $1', 2', \dots, k'$, with vertex i joined to vertex j if i and j belong to the same block of ρ. The representation of a partition by a diagram is not unique; for example the partition

$$\rho = \{\{1, 1', 3, 4', 5', 6\}, \{2, 2', 3', 4, 5, 6'\}\}$$

can be represented by the diagrams:

$\rho =$  or $\rho =$ 

If $\rho_1, \rho_2 \in A_k$, then the composition $\rho_1 \circ \rho_2$ is the partition obtained by placing ρ_1 above ρ_2 and identifying each vertex in the bottom row of ρ_1 with the corresponding vertex in the top row of ρ_2 and deleting any components of the resulting diagram which contains only elements from the middle row. The composition product makes A_k into an associative monoid with identity

$1 =$ 

Let z be an indeterminate and $R = \mathbb{Z}[z]$. The partition algebra $\mathcal{A}_k(z)$ is the R–module freely generated by A_k, equipped with the product

$$\rho_1\rho_2 = z^{\ell}\rho_1 \circ \rho_2, \qquad\qquad \text{for } \rho_1, \rho_2 \in A_k,$$

where ℓ is the number of blocks removed from the middle row in constructing the composition $\rho_1 \circ \rho_2$. Let $\mathcal{A}_{k-\frac{1}{2}}(z)$ denote the subalgebra of $\mathcal{A}_k(z)$ generated by $A_{k-\frac{1}{2}}$. A presentation for $\mathcal{A}_k(z)$ has been given by Halverson and Ram [HR] and East [Ea].

Theorem 2.1 (Theorem 1.11 of [HR])**.** *If $k = 1, 2, \ldots$, then the partition algebra $\mathcal{A}_k(z)$ is the unital associative R–algebra presented by the generators*

$$p_1, p_{1+\frac{1}{2}}, p_2, \ldots, p_k, s_1, s_2, \ldots, s_{k-1},$$

and the relations

- *(1) (Coxeter relations)*
  - *(i) $s_i^2 = 1$, for $i = 1, \ldots, k-1$.*
  - *(ii) $s_i s_j = s_j s_i$, if $j \neq i+1$.*
  - *(iii) $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, for $i = 1, \ldots, k-2$.*
- *(2) (Idempotent relations)*
  - *(i) $p_i^2 = zp_i$, for $i = 1, \ldots, k$.*
  - *(ii) $p_{i+\frac{1}{2}}^2 = p_{i+\frac{1}{2}}$, for $i = 1, \ldots, k-1$.*
  - *(iii) $s_i p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} s_i = p_{i+\frac{1}{2}}$, for $i = 1, \ldots, k-1$.*
  - *(iv) $s_i p_i p_{i+1} = p_i p_{i+1} s_i = p_i p_{i+1}$, for $i = 1, \ldots, k-1$.*
- *(3) (Commutation relations)*
  - *(i) $p_i p_j = p_j p_i$, for $i = 1, \ldots k$ and $j = 1, \ldots, k$.*
  - *(ii) $p_{i+\frac{1}{2}} p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} p_{i+\frac{1}{2}}$, for $i = 1, \ldots k-1$ and $j = 1, \ldots, k-1$.*
  - *(iii) $p_i p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} p_i$, for $j \neq i, i+1$.*
  - *(iv) $s_i p_j = p_j s_i$, for $j \neq i, i+1$.*
  - *(v) $s_i p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} s_i$, for $j \neq i-1, i+1$.*
  - *(vi) $s_i p_i s_i = p_{i+1}$, for $i = 1, \ldots, k-1$.*
  - *(vii) $s_i p_{i-\frac{1}{2}} s_i = s_{i-1} p_{i+\frac{1}{2}} s_{i-1}$, for $i = 2, \ldots, k-1$.*
- *(4) (Contraction relations)*
  - *(i) $p_{i+\frac{1}{2}} p_j p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, for $j = i, i+1$.*
  - *(ii) $p_i p_{j-\frac{1}{2}} p_i = p_i$, for $j = i, i+1$.*

The following identifications have been made in Theorem 2.1:

$s_i =$ 

and

$p_j =$ 

and

$p_{i+\frac{1}{2}} =$ 

We also recall the presentation for $\mathcal{A}_k(z)$ given in [En].

Theorem 2.2 (Theorem 4.1 of [En])**.** *If $k = 1, 2, \ldots,$ then $\mathcal{A}_k(z)$ is the unital associative algebra presented by the generators*

$$p_1, p_{1+\frac{1}{2}}, p_2, \ldots, p_k, \sigma_2, \sigma_{2+\frac{1}{2}}, \sigma_3, \ldots, \sigma_k,$$

and the relations:

- *(1) (Involutions)*
  - *(a) $\sigma^2_{i+\frac{1}{2}} = 1$, for $i = 2, \ldots, k-1$.*
  - *(b) $\sigma^2_{i+1} = 1$, for $i = 1, \ldots, k-1$.*
- *(2) (Braid–like relations)*
  - *(a) $\sigma_{i+1}\sigma_{j+\frac{1}{2}} = \sigma_{j+\frac{1}{2}}\sigma_{i+1}$, if $j \neq i+1$.*
  - *(b) $\sigma_i\sigma_j = \sigma_j\sigma_i$, if $j \neq i+1$.*
  - *(c) $\sigma_{i+\frac{1}{2}}\sigma_{j+\frac{1}{2}} = \sigma_{j+\frac{1}{2}}\sigma_{i+\frac{1}{2}}$, if $j \neq i+1$.*
  - *(d) $s_is_{i+1}s_i = s_{i+1}s_is_{i+1}$, for $i = 1, \ldots, k-2$, where*

    $$s_\ell = \begin{cases} \sigma_{\ell+1}, & \text{if } \ell = 1, \\ \sigma_{\ell+\frac{1}{2}}\sigma_{\ell+1}, & \text{if } \ell = 2, \ldots, k-1, \end{cases}$$

    are the Coxeter generators for the symmetric group.
- *(3) (Idempotent relations)*
  - *(a) $p_i^2 = zp_i$, for $i = 1, \ldots, k$.*
  - *(b) $p^2_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, for $i = 1, \ldots, k-1$.*
  - *(c) $\sigma_{i+1}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}\sigma_{i+1} = p_{i+\frac{1}{2}}$, for $i = 1, \ldots, k-1$.*
  - *(d) $\sigma_{i+\frac{1}{2}}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}\sigma_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, for $i = 1, \ldots, k-1$.*
  - *(e) $\sigma_{i+\frac{1}{2}}p_ip_{i+1} = \sigma_{i+1}p_ip_{i+1}$, for $i = 1, \ldots, k-1$.*
  - *(f) $p_ip_{i+1}\sigma_{i+\frac{1}{2}} = p_ip_{i+1}\sigma_{i+1}$, for $i = 1, \ldots, k-1$.*
- *(4) (Commutation relations)*
  - *(a) $p_ip_j = p_jp_i$, for $i, j = 1, \ldots, k$.*
  - *(b) $p_{i+\frac{1}{2}}p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}}p_{i+\frac{1}{2}}$, for $i, j = 1, \ldots, k-1$.*
  - *(c) $p_{i+\frac{1}{2}}p_j = p_jp_{i+\frac{1}{2}}$, for $j \neq i, i+1$.*
  - *(d) $\sigma_ip_j = p_j\sigma_i$ if $j \neq i-1, i$.*
  - *(e) $\sigma_ip_{j+\frac{1}{2}} = p_{j+\frac{1}{2}}\sigma_i$, if $j \neq i$.*
  - *(f) $\sigma_{i+\frac{1}{2}}p_j = p_j\sigma_{i+\frac{1}{2}}$, if $j \neq i, i+1$.*
  - *(g) $\sigma_{i+\frac{1}{2}}p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}}\sigma_{i+\frac{1}{2}}$, if $j \neq i-1$.*
  - *(h) $\sigma_{i+\frac{1}{2}}p_i\sigma_{i+\frac{1}{2}} = \sigma_{i+1}p_{i+1}\sigma_{i+1}$, for $i = 1, \ldots, k-1$.*
  - *(i) $\sigma_{i+\frac{1}{2}}p_{i-\frac{1}{2}}\sigma_{i+\frac{1}{2}} = \sigma_ip_{i+\frac{1}{2}}\sigma_i$, for $i = 2, \ldots, k-1$.*
- *(5) (Contraction relations)*
  - *(a) $p_{i+\frac{1}{2}}p_jp_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, for $j = i, i+1$.*
  - *(b) $p_ip_{j-\frac{1}{2}}p_i = p_i$, for $j = i, i+1$.*

It will be useful to note that R-linear map $* : \mathcal{A}_k(z) \to \mathcal{A}_k(z)$ defined on generators by

$$u^* = u^{-1} \qquad\qquad (\text{for } u \in \langle \sigma_2, \sigma_{2+\frac{1}{2}}, \ldots, \sigma_k \rangle)$$

and

$$p_i^* = p_i \qquad (\text{for } i = 1, \ldots, k) \qquad \text{and} \qquad p^*_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} \qquad (\text{for } j = 1, \ldots, k-1),$$

is an algebra anti-involution of $\mathcal{A}_k(z)$. Restricting the map $*$ from $\mathcal{A}_k(z)$ to $\mathcal{A}_{k-\frac{1}{2}}(z)$, gives an algebra anti involution of $\mathcal{A}_{k-\frac{1}{2}}(z)$ which we also denote by $*$.

2.3. **Jucys–Murphy elements.** Jucys–Murphy elements for the partition algebras were defined in the diagram basis by Halverson and Ram [HR]. The following recursions for Jucys–Murphy elements in $\mathcal{A}_k(z)$ and $\mathcal{A}_{k+\frac{1}{2}}(z)$, which have been obtained in [En], are equivalent to the definition of Jucys–Murphy elements given in [HR].

Let $(\sigma_i : i = 1, 2, \dots)$ and $(L_i : i = 0, 1, \dots)$ be given by

$$L_0 = 0, \qquad L_1 = p_1, \qquad \sigma_1 = 1, \qquad \text{and,} \qquad \sigma_2 = s_1,$$

and, for $i = 1, 2, \dots,$

$$L_{i+1} = -s_iL_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_is_i + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + s_iL_is_i + \sigma_{i+1}, \tag{2.3}$$

where, for $i = 2, 3, \dots,$

$$\begin{aligned}\sigma_{i+1} = s_{i-1}s_i\sigma_is_is_{i-1} + s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}s_i + p_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}\\ - s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} - p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_i.\end{aligned} \tag{2.4}$$

Define $(\sigma_{i+\frac{1}{2}} : i = 1, 2, \dots)$ and $(L_{i+\frac{1}{2}} : i = 0, 1, \dots)$ by

$$L_{\frac{1}{2}} = 0, \qquad \sigma_{\frac{1}{2}} = 1, \qquad \text{and,} \qquad \sigma_{1+\frac{1}{2}} = 1,$$

and, for $i = 1, 2, \dots,$

$$L_{i+\frac{1}{2}} = -L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i + p_{i+\frac{1}{2}}L_ip_ip_{i+\frac{1}{2}} + s_iL_{i-\frac{1}{2}}s_i + \sigma_{i+\frac{1}{2}}, \tag{2.5}$$

where, for $i = 2, 3, \dots,$

$$\begin{aligned}\sigma_{i+\frac{1}{2}} = s_{i-1}s_i\sigma_{i-\frac{1}{2}}s_is_{i-1} + p_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}s_i + s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}\\ - p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} - s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_i.\end{aligned} \tag{2.6}$$

Remark 2.3. The transition from the algebra presentation given in Theorem 2.1 to the presentation in Theorem 2.2 is defined by the equations (2.3)–(2.6).

We collate the following facts from §3 of [En] for later reference.

Proposition 2.4. *For $i = 1, 2, \dots,$ the following statements hold:*

(1) $\sigma_{i+\frac{1}{2}} \in \mathcal{A}_{i+\frac{1}{2}}(z)$ *and* $\sigma_{i+1} \in \mathcal{A}_{i+1}(z)$.
(2) $(\sigma_{i+\frac{1}{2}})^* = \sigma_{i+\frac{1}{2}}$ *and* $(\sigma_{i+1})^* = \sigma_{i+1}$.
(3) $L_{i+\frac{1}{2}} \in \mathcal{A}_{i+\frac{1}{2}}(z)$ *and* $L_{i+1} \in \mathcal{A}_{i+1}(z)$.
(4) $(L_{i+\frac{1}{2}})^* = L_{i+\frac{1}{2}}$ *and* $(L_{i+1})^* = \sigma_{i+1}$.
(5) $s_{i+1}\sigma_{i+1}p_{i+\frac{3}{2}} = p_{i+\frac{1}{2}}s_{i+1}\sigma_{i+1}$.
(6) $\sigma_{i+1}p_ip_{i+\frac{1}{2}} = s_iL_ip_{i+\frac{1}{2}}$.
(7) $\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} = L_ip_{i+\frac{1}{2}}$.
(8) $\sigma_{i+\frac{1}{2}}\sigma_{i+1} = \sigma_{i+1}\sigma_{i+\frac{1}{2}} = s_i$, *and* $\sigma^2_{i+\frac{1}{2}} = \sigma^2_{i+1} = 1$.
(9) L_{i+1} *commutes with* $\mathcal{A}_{i+\frac{1}{2}}(z)$, *and* σ_{i+1} *commutes with* $\mathcal{A}_{i-\frac{1}{2}}(z)$.
(10) $L_{i+\frac{1}{2}}$ *commutes with* $\mathcal{A}_i(z)$, *and* $\sigma_{i+\frac{1}{2}}$ *commutes with* $\mathcal{A}_{i-1}(z)$.
(11) $(L_{i+\frac{1}{2}} + L_{i+1})p_{i+1} = p_{i+1}(L_{i+\frac{1}{2}} + L_{i+1}) = zp_{i+1}$.
(12) $(L_i + L_{i+\frac{1}{2}})p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}(L_i + L_{i+\frac{1}{2}}) = zp_{i+\frac{1}{2}}$.
(13) The element $z_i = L_{\frac{1}{2}} + L_1 + L_{1+\frac{1}{2}} + \cdots + L_i$ *is central in* $\mathcal{A}_i(z)$.
(14) The element $z_{i+\frac{1}{2}} = L_{\frac{1}{2}} + L_1 + L_{1+\frac{1}{2}} + \cdots + L_{i+\frac{1}{2}}$ *is central in* $\mathcal{A}_{i+\frac{1}{2}}(z)$.
(15) $p_{i+1}\sigma_{i+1}p_{i+1} = L_ip_{i+1}$.
(16) $p_{i+1}\sigma_{i+\frac{1}{2}}p_{i+1} = (z - L_{i-\frac{1}{2}})p_{i+1}$.
(17) $p_{i+\frac{3}{2}}\sigma_{i+1}p_{i+\frac{3}{2}} = p_{i+\frac{1}{2}}p_{i+\frac{3}{2}}$.

The next statement gives recursions for the Jucys–Murphy elements in terms of the presentation Theorem 2.2.

Proposition 2.5. *For $i = 1, 2, \ldots,$ the following statements hold:*
(1) $\sigma_{i+1}L_{i+1} - L_i\sigma_{i+1} = -L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}p_{i+1} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + 1.$
(2) $\sigma_{i+\frac{1}{2}}L_{i+\frac{1}{2}} - L_{i-\frac{1}{2}}\sigma_{i+\frac{1}{2}} = -p_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i + (z - L_{i-\frac{1}{2}})p_{i+\frac{1}{2}} + 1.$
(3) $L_{i+1} = -p_{i+1}p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}p_{i+1} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + \sigma_{i+1}L_i\sigma_{i+1} + \sigma_{i+1}.$
(4) $L_{i+\frac{1}{2}} = -p_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}p_i + (z - L_{i-\frac{1}{2}})p_{i+\frac{1}{2}} + \sigma_{i+\frac{1}{2}}L_i\sigma_{i+\frac{1}{2}} + \sigma_{i+\frac{1}{2}}.$

Proof. (1) The definition (2.3), and the relations

$$\sigma_{i+1}s_i = s_i\sigma_{i+1} = \sigma_{i+\frac{1}{2}} \qquad \text{and} \qquad \sigma_{i+1}^2 = 1,$$

imply that

$$s_iL_{i+1}s_i = -L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + L_i + \sigma_{i+1},$$

and

$$s_iL_{i+1}\sigma_{i+\frac{1}{2}} = -L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i\sigma_{i+1} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + L_i\sigma_{i+1} + 1.$$

Since L_{i+1} commutes with $\sigma_{i+\frac{1}{2}} \in \mathcal{A}_{i+\frac{1}{2}}(x)$,

$$\sigma_{i+1}L_{i+1} = -L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i\sigma_{i+1} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + L_i\sigma_{i+1} + 1,$$

and

$$\sigma_{i+1}L_{i+1} - L_i\sigma_{i+1} = -L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i\sigma_{i+1} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + 1.$$

Making the substitution $p_{i+\frac{1}{2}}p_{i+1}\sigma_{i+1} = p_{i+\frac{1}{2}}L_i$, or $p_{i+\frac{1}{2}}p_{i+1} = p_{i+\frac{1}{2}}L_i\sigma_{i+1}$ in the last expression gives the required statement.
(2) Since

$$L_{i+\frac{1}{2}} = -L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i + (z - L_{i-\frac{1}{2}})p_{i+\frac{1}{2}} + s_iL_{i-\frac{1}{2}}s_i + \sigma_{i+\frac{1}{2}},$$

the relations $\sigma_{i+1}s_i = s_i\sigma_{i+1} = \sigma_{i+\frac{1}{2}}$ and $\sigma_{i+1}^2 = 1$ imply that

$$\sigma_{i+\frac{1}{2}}L_{i+\frac{1}{2}} = -\sigma_{i+\frac{1}{2}}L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i + (z - L_{i-\frac{1}{2}})p_{i+\frac{1}{2}} + \sigma_{i+1}L_{i-\frac{1}{2}}s_i + 1.$$

Since σ_{i+1} commutes with $L_{i-\frac{1}{2}} \in \mathcal{A}_{i-\frac{1}{2}}(x)$,

$$\sigma_{i+\frac{1}{2}}L_{i+\frac{1}{2}} - L_{i-\frac{1}{2}}\sigma_{i+\frac{1}{2}} = -\sigma_{i+\frac{1}{2}}L_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i + (z - L_{i-\frac{1}{2}})p_{i+\frac{1}{2}} + 1.$$

Applying the relation $p_{i+\frac{1}{2}}L_i\sigma_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_i$ from Proposition 2.4 to the right hand side of the last equality, gives the required statement.
(3) Using item (7) of Proposition 2.4 and the relation $\sigma_{i+1}^2 = 1$, we obtain $\sigma_{i+1}L_ip_{i+\frac{1}{2}} = p_{i+1}p_{i+\frac{1}{2}}$. Since $\sigma_{i+1}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, the statement follows from item (1).
(4) We proceed as in (3), using the relation $p_{i+\frac{1}{2}}L_i\sigma_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_i$. ☐

3. A MURPHY BASIS

Murphy–type bases for the partition algebras have been given in [En1] (see Theorems 3.2 and 3.3). In this section, we re–write the bases of [En1] and use the new definition of Murphy–type bases to show that the Jucys–Murphy elements (2.3) and (2.5) act triangularly on the partition algebras.

For $k = 0, 1, \ldots,$ let

$$\hat{A}_k = \hat{A}_{k+\frac{1}{2}} = \{(\lambda, \ell) \mid \lambda \vdash (k - \ell), \text{ for } \ell = 0, 1, \ldots, k\}.$$

If $(\lambda,\ell),(\mu,m)\in\hat{A}_k$, write $(\lambda,\ell)\trianglerighteq(\mu,m)$ if either (i) $l=m$ and $\lambda\trianglerighteq\mu$, or (ii) $\ell>m$.

Following §2 of [HR], build a graph $\hat{A}$ with

(1) vertices on level k: $\hat{A}_k$,
(2) vertices on level $k+\frac{1}{2}$: $\hat{A}_{k+\frac{1}{2}}=\hat{A}_k$,
(3) an edge $(\lambda,\ell)\to(\mu,m)$ in $\hat{A}$, for $(\lambda,\ell)\in\hat{A}_k$, $(\mu,m)\in\hat{A}_{k+\frac{1}{2}}$, if $\lambda=\mu$, or if λ is obtained from μ by removing a node,
(4) an edge $(\lambda,\ell)\to(\mu,m)$ in $\hat{A}$, for $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$, $(\mu,m)\in\hat{A}_{k+1}$, if $\lambda=\mu$, or if λ is obtained from μ by adding a node.

Definition 3.1. Let $j\in\frac{1}{2}\mathbb{Z}_{\geqslant0}$. A path to level j in the graph $\hat{A}$ is a sequence

$$\mathfrak{t}=\big((\lambda^{(0)},\ell_0),(\lambda^{(\frac{1}{2})},\ell_{\frac{1}{2}}),(\lambda^{(1)},\ell_1),\ldots,(\lambda^{(j)},\ell_j)\big),\qquad\text{where}\tag{3.1}$$

$$(\lambda^{(0)},\ell_0)=(\emptyset,0)\qquad\text{and}\qquad(\lambda^{(r)},\ell_r)\to(\lambda^{(r+\frac{1}{2})},\ell_{r+\frac{1}{2}}),\qquad\text{for } r=0,\tfrac{1}{2},1,\ldots,j-\tfrac{1}{2}.$$

We say that $\mathfrak{t}$ is a path of shape (λ,ℓ), and write $(\lambda,\ell)=\mathrm{Shape}(\mathfrak{t})$.

If $(\lambda,\ell)\in\hat{A}_k$, let

$$A_k^{(\lambda,\ell)}=\big\{\mathfrak{t}\mid\mathfrak{t}\text{ is a path to level }k\text{, and }\mathrm{Shape}(\mathfrak{t})=(\lambda,\ell)\big\},\qquad\text{and}$$
$$A_{k+\frac{1}{2}}^{(\lambda,\ell)}=\big\{\mathfrak{t}\mid\mathfrak{t}\text{ is a path to level }k+\tfrac{1}{2}\text{, and }\mathrm{Shape}(\mathfrak{t})=(\lambda,\ell)\big\}.$$

The definitions of the sets $\hat{A}_k^{(\lambda,\ell)}$ and $\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$, for $(\lambda,\ell)\in\hat{A}_k$, are illustrated in the diagram (3.2) where the first few levels of the graph $\hat{A}$ are given.



(3.2)

If $\mathfrak{t}$ is a path of the form (3.1), we will generally write

$$\mathfrak{t}=\big(\lambda^{(0)},\lambda^{(\frac{1}{2})},\lambda^{(1)},\ldots,\lambda^{(j)}\big),$$

and, if $r=0,\frac{1}{2},1,\ldots,j-\frac{1}{2}$, define the truncation of $\mathfrak{t}$ to level r to be the path

$$\mathfrak{t}|_r=\big(\lambda^{(0)},\lambda^{(\frac{1}{2})},\lambda^{(1)},\ldots,\lambda^{(r)}\big).$$

We can now define the Murphy bases for partition algebras. For $i = 1, 2, \dots,$ let

$$p_i^{(\ell)} = \underbrace{p_{i-\ell+1}p_{i-\ell+2}\cdots p_i}_{\ell \text{ factors}} \qquad \text{and} \qquad p_{i+\frac{1}{2}}^{(\ell)} = \underbrace{p_{i-\ell+\frac{3}{2}}p_{i-\ell+\frac{5}{2}}\cdots p_{i+\frac{1}{2}}}_{\ell \text{ factors}} \qquad \text{if } \ell \leqslant i$$

and let

$$p_i^{(\ell)} = 0 \qquad \text{and} \qquad p_{i+\frac{1}{2}}^{(\ell)} = 0 \qquad \text{if } \ell > i.$$

For $i = 0, 1, \dots,$ and $(\lambda, \ell) \in \hat{A}_i$, let

$$c_\lambda = \sum_{v \in \mathfrak{S}_\lambda} v$$

and

$$x_{(\lambda,\ell)}^{(i)} = c_\lambda p_i^{(\ell)} \in \mathcal{A}_i \qquad \text{and} \qquad x_{(\lambda,\ell)}^{(i+\frac{1}{2})} = c_\lambda p_{i+\frac{1}{2}}^{(\ell)} \in \mathcal{A}_{i+\frac{1}{2}}.$$

Define the two sided ideals

$$\mathcal{A}_i^{\rhd(\lambda,\ell)} = \sum_{(\mu,m)\rhd(\lambda,\ell)} \mathcal{A}_i x_{(\mu,m)}^{(i)} \mathcal{A}_i \qquad \text{and} \qquad \mathcal{A}_{i+\frac{1}{2}}^{\rhd(\lambda,\ell)} = \sum_{(\mu,m)\rhd(\lambda,\ell)} \mathcal{A}_{i+\frac{1}{2}} x_{(\mu,m)}^{(i+\frac{1}{2})} \mathcal{A}_{i+\frac{1}{2}}$$

where each sum is taken over $(\mu, m) \in \hat{A}_i$ such that $(\mu, m) \rhd (\lambda, \ell)$. For $(\lambda, \ell) \in \hat{A}_i$, define the right $\mathcal{A}_i$–module

$$A_i^{(\lambda,\ell)} = \left\{ x_{(\lambda,\ell)}^{(i)} p + \mathcal{A}_i^{\rhd(\lambda,\ell)} \,\middle|\, p \in \mathcal{A}_i \right\} \subseteq \mathcal{A}_i / \mathcal{A}_i^{\rhd(\lambda,\ell)}$$

and the right $\mathcal{A}_{i+\frac{1}{2}}$–module

$$A_{i+\frac{1}{2}}^{(\lambda,\ell)} = \left\{ x_{(\lambda,\ell)}^{(i+\frac{1}{2})} p + \mathcal{A}_{i+\frac{1}{2}}^{\rhd(\lambda,\ell)} \,\middle|\, p \in \mathcal{A}_{i+\frac{1}{2}} \right\} \subseteq \mathcal{A}_{i+\frac{1}{2}} / \mathcal{A}_{i+\frac{1}{2}}^{\rhd(\lambda,\ell)}$$

If $\lambda \vdash (i-1)$ and $\mu \vdash i$, such that $\mu = \lambda \cup \{(j, \mu_j)\}$, let $a_j = \sum_{r=1}^{j} \mu_j$ and define

$$\bar{u}_{\lambda\to\mu}^{(i)} = w_{i,a_j} \sum_{r=0}^{\lambda_j} w_{a_j, a_j - r} \qquad \text{and} \qquad \bar{d}_{\lambda\to\mu}^{(i)} = w_{a_j, i},$$

so that

$$c_\lambda^{(i-1)} \bar{u}_{\lambda\to\mu}^{(i)} = \big(\bar{d}_{\lambda\to\mu}^{(i)}\big)^* c_\mu^{(i)}.$$

If $(\lambda, \ell) \in \hat{A}_{i-\frac{1}{2}}$ and $(\mu, m) \in \hat{A}_i$ and $(\lambda, \ell) \to (\mu, m)$, let

$$a_{(\lambda,\ell)\to(\mu,m)}^{(i)} = \begin{cases} \bar{d}_{\lambda\to\mu}^{(i-m)} p_{i-1}^{(m)}, & \text{if } \ell = m, \\ p_{i-1}^{(m-1)}, & \text{if } \ell = m-1. \end{cases}$$

Similarly, if $(\lambda, \ell) \in \hat{A}_i$ and $(\mu, m) \in \hat{A}_{i+\frac{1}{2}}$ and $(\lambda, \ell) \to (\mu, m)$, let

$$a_{(\lambda,\ell)\to(\mu,m)}^{(i+\frac{1}{2})} = \begin{cases} p_{i-\frac{1}{2}}^{(m)}, & \text{if } m = n, \\ p_{i-\frac{1}{2}}^{(m-1)} \bar{u}_{\lambda\to\mu}^{(i-m+1)}, & \text{if } \ell = m-1. \end{cases}$$

Generally, we write $a_{\lambda\to\mu}^{(i)}$ for $a_{(\lambda,\ell)\to(\mu,m)}^{(i)}$ and $a_{\lambda\to\mu}^{(i+\frac{1}{2})}$ for $a_{(\lambda,\ell)\to(\mu,m)}^{(i+\frac{1}{2})}$.

Let $k \in \mathbb{Z}_{\geqslant 0}$ and $(\lambda, \ell) \in \hat{A}_k$. For $\mathfrak{t} = (\lambda^{(0)}, \lambda^{(\frac{1}{2})}, \dots, \lambda^{(k)}) \in \hat{A}_k^{(\lambda,\ell)}$, let

$$a_{\mathfrak{t}}^{(k)} = a_{\lambda^{(k-\frac{1}{2})}\to\lambda^{(k)}}^{(k)} a_{\lambda^{(k-1)}\to\lambda^{(k-\frac{1}{2})}}^{(k-\frac{1}{2})} \cdots a_{\lambda^{(0)}\to\lambda^{(\frac{1}{2})}}^{(\frac{1}{2})},$$

and, for $\mathfrak{s} = (\lambda^{(0)}, \lambda^{(\frac{1}{2})}, \dots, \lambda^{(k+\frac{1}{2})}) \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, let

$$a_{\mathfrak{s}}^{(k+\frac{1}{2})} = a^{(k+\frac{1}{2})}_{\lambda^{(k)} \to \lambda^{(k+\frac{1}{2})}} a^{(k)}_{\lambda^{(k-\frac{1}{2})} \to \lambda^{(k)}} \cdots a^{(\frac{1}{2})}_{\lambda^{(0)} \to \lambda^{(\frac{1}{2})}}.$$

The two statements below are Theorem 5.10 of [En1] applied to the algebras $\mathcal{A}_k$ and $\mathcal{A}_{k+\frac{1}{2}}$ respectively. For brevity we have written $a_{\mathfrak{t}} = a_{\mathfrak{t}}^{(k)}$ in Theorem 3.2, and $a_{\mathfrak{t}} = a_{\mathfrak{t}}^{(k+\frac{1}{2})}$ in Theorem 3.3.

Theorem 3.2. *If $k = 1, 2, \dots,$ then the set*

$$\mathscr{A}_k = \left\{ a_{\mathfrak{s}}^* x^{(k)}_{(\lambda,\ell)} a_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_k,\ (\lambda,\ell) \in \hat{A}_k,\ \text{and } \ell = 0, 1, \dots, k \right\} \tag{3.3}$$

is an R–basis for $\mathcal{A}_k$. Moreover, the following statements hold:

(1) If $(\lambda,\ell) \in \hat{A}_k$, $\mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_k$, and $p \in \mathcal{A}_k$, there exist $r_{\mathfrak{u}} \in R$, for $\mathfrak{u} \in \hat{A}^{(\lambda,\ell)}_k$, such that

$$a_{\mathfrak{s}}^* x^{(k)}_{(\lambda,\ell)} a_{\mathfrak{t}} p \equiv \sum_{\mathfrak{u} \in \hat{A}^{(\lambda,\ell)}_k} r_{\mathfrak{u}} a_{\mathfrak{s}}^* x^{(k)}_{(\lambda,\ell)} a_{\mathfrak{u}} \mod A^{\rhd(\lambda,\ell)}_k \qquad \text{for all } \mathfrak{s} \in \hat{A}^{(\lambda,\ell)}_k,$$

where $\mathcal{A}^{\rhd(\lambda,\ell)}_k$ is the R–module freely generated by

$$\left\{ a_{\mathfrak{s}}^* x^{(k)}_{(\mu,m)} a_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{A}^{(\mu,m)}_k,\ (\mu,m) \in \hat{A}_k,\ \text{and } (\mu,m) \rhd (\lambda,\ell) \right\}.$$

(2) If $(\lambda,\ell) \in \hat{A}_k$, and $\mathfrak{s}, \mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_k$, then $* : a_{\mathfrak{s}}^* x^{(k)}_{(\lambda,\ell)} a_{\mathfrak{t}} \mapsto a_{\mathfrak{t}}^* x^{(k)}_{(\lambda,\ell)} a_{\mathfrak{s}}$.

Theorem 3.3. *If $k = 1, 2, \dots,$ then the set*

$$\mathscr{A}_{k+\frac{1}{2}} = \left\{ a_{\mathfrak{s}}^* x^{(k+\frac{1}{2})}_{(\lambda,\ell)} a_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}},\ (\lambda,\ell) \in \hat{A}_{k+\frac{1}{2}},\ \text{and } \ell = 0, 1, \dots, k \right\} \tag{3.4}$$

is an R–basis for $\mathcal{A}_{k+\frac{1}{2}}$. Moreover, the following statements hold:

(1) If $(\lambda,\ell) \in \hat{A}_{k+\frac{1}{2}}$, $\mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, and $p \in \mathcal{A}_{k+\frac{1}{2}}$, there exist $r_{\mathfrak{u}} \in R$, for $\mathfrak{u} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, such that

$$a_{\mathfrak{s}}^* x^{(k)}_{(\lambda,\ell)} a_{\mathfrak{t}} p \equiv \sum_{\mathfrak{u} \in \hat{A}^{(\lambda,\ell)}_k} r_{\mathfrak{u}} a_{\mathfrak{s}}^* x^{(k+\frac{1}{2})}_{(\lambda,\ell)} a_{\mathfrak{u}} \mod A^{\rhd(\lambda,\ell)}_{k+\frac{1}{2}} \qquad \text{for all } \mathfrak{s} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}},$$

where $\mathcal{A}^{\rhd(\lambda,\ell)}_{k+\frac{1}{2}}$ is the R–module freely generated by

$$\left\{ a_{\mathfrak{s}}^* x^{(k+\frac{1}{2})}_{(\mu,m)} a_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{A}^{(\mu,m)}_{k+\frac{1}{2}},\ (\mu,m) \in \hat{A}_{k+\frac{1}{2}},\ \text{and } (\mu,m) \rhd (\lambda,\ell) \right\}.$$

(2) If $(\lambda,\ell) \in \hat{A}_{k+\frac{1}{2}}$, and $\mathfrak{s}, \mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, then $* : a_{\mathfrak{s}}^* x^{(k+\frac{1}{2})}_{(\lambda,\ell)} a_{\mathfrak{t}} \mapsto a_{\mathfrak{t}}^* x^{(k+\frac{1}{2})}_{(\lambda,\ell)} a_{\mathfrak{s}}$.

The next statement gives a filtration of the cell modules for $\mathcal{A}_i$ by $\mathcal{A}_{i-\frac{1}{2}}$–modules and a filtration of the cell modules for $\mathcal{A}_{i+\frac{1}{2}}$ by $\mathcal{A}_i$–modules.

Proposition 3.4 (Proposition 4.2 of [En1])**.** *(1) Let $(\mu,m) \in \hat{A}_i$ and $\{(\lambda^{(1)},\ell_1), \dots, (\lambda^{(t)},\ell_t)\}$ be an indexing of the set*

$$\{(\lambda,\ell) \in \hat{A}_{i-\frac{1}{2}} \mid (\lambda,\ell) \to (\mu,m)\}$$

such that $(\lambda^{(r)},\ell_r) \rhd (\lambda^{(s)},\ell_s)$ whenever $s > r$. For $j = 1, \dots, t$, let

$$N_j = \sum_{\substack{\mathfrak{s} \in \hat{A}^{(\mu,m)}_i \\ \mathrm{Shape}(\mathfrak{s}|_{i-\frac{1}{2}}) \trianglerighteq (\lambda^{(j)},\ell_j)}} \left(x^{(i)}_{(\mu,m)} + A^{\rhd(\mu,m)}_i \right) a^{(i)}_{\mathfrak{s}} A_{i-\frac{1}{2}}.$$

Then

$$\{0\} = N_0 \subseteq N_1 \subseteq \cdots \subseteq N_p = A_i^{(\mu,m)}$$

is a filtration by right $A_{i-\frac{1}{2}}$–modules and, for $j = 1, \dots, t$, the R–linear map

$$\begin{aligned} A_{i-\frac{1}{2}}^{(\lambda^{(j)},\ell_j)} &\longrightarrow N_j/N_{j-1} \\ \big(x^{(i-\frac{1}{2})}_{(\lambda^{(j)},\ell_j)} + A^{\rhd(\mu,m)}_{i-\frac{1}{2}}\big) a^{(i-\frac{1}{2})}_{\mathfrak{u}} &\longmapsto \big(x^{(i)}_{(\mu,m)} + A_i^{\rhd(\mu,m)}\big) a^{(i)}_{\mathfrak{t}} + N_{j-1}, \end{aligned} \tag{3.5}$$

for $\mathfrak{u} \in \hat{A}_{i-\frac{1}{2}}^{(\lambda^{(j)},\ell_j)}$ and $\mathfrak{t} \in \hat{A}_i^{(\mu,m)}$ such that $\mathfrak{t}|_{i-\frac{1}{2}} = \mathfrak{u}$, is an isomorphism of right $A_{i-\frac{1}{2}}$–modules.
(2) Let $(\mu, m) \in \hat{A}_{i+\frac{1}{2}}$ and $\{(\lambda^{(1)}, \ell_1), \dots, (\lambda^{(t)}, \ell_t)\}$ be an indexing of the set

$$\{(\lambda, \ell) \in \hat{A}_i \mid (\lambda, \ell) \to (\mu, m)\}$$

such that $(\lambda_r, \ell_r) \rhd (\lambda_s, \ell_s)$ whenever $s > r$. For $j = 1, \dots, t$, let

$$N_j = \sum_{\substack{\mathfrak{s} \in \hat{A}^{(\mu,m)}_{i+\frac{1}{2}} \\ \mathrm{Shape}(\mathfrak{s}|_i) \unrhd (\lambda^{(j)},\ell_j)}} \big(x^{(i+\frac{1}{2})}_{(\mu,m)} + A^{\rhd(\mu,m)}_{i+\frac{1}{2}}\big) a^{(i+\frac{1}{2})}_{\mathfrak{s}} A_i.$$

Then

$$\{0\} = N_0 \subseteq N_1 \subseteq \cdots \subseteq N_p = A_{i+\frac{1}{2}}^{(\mu,m)}$$

is a filtration by right A_i–modules and, for $j = 1, \dots, p$, the R–linear map

$$\begin{aligned} A_i^{(\lambda_j,\ell_j)} &\longrightarrow N_j/N_{j-1} \\ \big(x^{(i)}_{(\lambda_j,\ell_j)} + A_i^{\rhd(\mu,m)}\big) a_{\mathfrak{u}} &\longmapsto \big(x^{(i+\frac{1}{2})}_{(\mu,m)} + A^{\rhd(\mu,m)}_{i+\frac{1}{2}}\big) a^{(i+\frac{1}{2})}_{\mathfrak{t}} + N_{j-1}, \end{aligned} \tag{3.6}$$

for $\mathfrak{u} \in \hat{A}_i^{(\lambda_j,\ell_j)}$ and $\mathfrak{t} \in \hat{A}^{(\mu,m)}_{i+\frac{1}{2}}$ such that $\mathfrak{t}|_i = \mathfrak{u}$, is an isomorphism of right A_i–modules.

Corollary 3.5. *(1) Let $(\mu, m) \in \hat{A}_i$ and $(\lambda, \ell) \in \hat{A}_{i-\frac{1}{2}}$ such that $(\lambda, \ell) \to (\mu, m)$. If $p \in \mathcal{A}_{i-\frac{1}{2}}$ and $x^{(i-\frac{1}{2})}_{(\lambda,\ell)} p \in \mathcal{A}^{\rhd(\lambda,\ell)}_{i-\frac{1}{2}}$, then there exist $r_{\mathfrak{s}} \in R$, for $\mathfrak{s} \in \hat{A}_i^{(\mu,m)}$, such that*

$$x^{(i)}_{(\mu,m)} a^{(i)}_{\lambda\to\mu} p \equiv \sum_{\mathfrak{s} \in \hat{A}_i^{(\mu,m)}} r_{\mathfrak{s}} x^{(i)}_{(\mu,m)} a^{(i)}_{\mathfrak{s}} \mod \mathcal{A}_i^{\rhd(\mu,m)},$$

where $r_{\mathfrak{s}} = 0$ unless $\mathrm{Shape}(\mathfrak{s}|_{i-\frac{1}{2}}) \rhd (\lambda, \ell)$.
(2) Let $(\mu, m) \in \hat{A}_{i+\frac{1}{2}}$ and $(\lambda, \ell) \in \hat{A}_i$ such that $(\lambda, \ell) \to (\mu, m)$. If $p \in \mathcal{A}_i$ and $x^{(i)}_{(\lambda,\ell)} p \in \mathcal{A}_i^{\rhd(\lambda,\ell)}$, then there exist $r_{\mathfrak{s}} \in R$, for $\mathfrak{s} \in \hat{A}^{(\mu,m)}_{i+\frac{1}{2}}$, such that

$$x^{(i+\frac{1}{2})}_{(\mu,m)} a^{(i+\frac{1}{2})}_{\lambda\to\mu} p \equiv \sum_{\mathfrak{s} \in \hat{A}^{(\mu,m)}_{i+\frac{1}{2}}} r_{\mathfrak{s}} x^{(i+\frac{1}{2})}_{(\mu,m)} a^{(i+\frac{1}{2})}_{\mathfrak{s}} \mod \mathcal{A}^{\rhd(\mu,m)}_{i+\frac{1}{2}},$$

where $r_{\mathfrak{s}} = 0$ unless $\mathrm{Shape}(\mathfrak{s}|_i) \rhd (\lambda, \ell)$.

Proof. (1) Observe that there exists $b^{(i)}_{\lambda\to\mu} \in \mathcal{A}_i$ such that

$$x^{(i)}_{(\mu,m)} a^{(i)}_{\lambda\to\mu} = b^{(i)}_{\lambda\to\mu} x^{(i-\frac{1}{2})}_{(\lambda,\ell)}$$

and then use (1) of Proposition 3.4 to see that if $p \in \mathcal{A}_{i-\frac{1}{2}}$ and $x^{(i-\frac{1}{2})}_{(\lambda,\ell)} p \in \mathcal{A}^{\triangleright(\lambda,\ell)}_{i-\frac{1}{2}}$, then

$$b^{(i)}_{\lambda\to\mu} x^{(i-\frac{1}{2})}_{(\lambda,\ell)} p \equiv \sum_{\mathfrak{s}\in\hat{A}^{(\mu,m)}_i} r_{\mathfrak{s}} x^{(i)}_{(\mu,m)} a^{(i)}_{\mathfrak{s}} \mod \mathcal{A}^{\triangleright(\mu,m)}_i,$$

where $r_{\mathfrak{s}} = 0$ unless $\mathrm{Shape}(\mathfrak{s}|_{i-\frac{1}{2}}) \triangleright (\lambda,\ell)$. The proof of (2) is similar. □

The sets $\hat{A}^{(\lambda,\ell)}_i$ and $\hat{A}^{(\lambda,\ell)}_{i+\frac{1}{2}}$ are partially ordered by the following *reverse lexicographic order* given in §2.7 of [GG1].

Definition 3.6. Let $i \in \frac{1}{2}\mathbb{Z}$ and $\mathfrak{s} = (\mathfrak{s}_0, \mathfrak{s}_{\frac{1}{2}}, \ldots, \mathfrak{s}_i)$ and $\mathfrak{t} = (\mathfrak{t}_0, \mathfrak{t}_{\frac{1}{2}}, \ldots, \mathfrak{t}_i)$ be two paths in $\hat{A}$. We write $\mathfrak{s} \succcurlyeq \mathfrak{t}$ if $\mathfrak{s} = \mathfrak{t}$, or if for the last index $j \in \frac{1}{2}\mathbb{Z}$ such that $\mathfrak{s}^{(j)} \neq \mathfrak{t}^{(j)}$, we have $\mathfrak{s}_j \trianglerighteq \mathfrak{t}_j$ in $\hat{A}_j$. Let $\mathfrak{s} \succ \mathfrak{t}$ denote the fact that $\mathfrak{s} \succcurlyeq \mathfrak{t}$ and $\mathfrak{s} \neq \mathfrak{t}$; write $\mathfrak{s} \overset{j}{\succ} \mathfrak{t}$ if $\mathfrak{s} \succ \mathfrak{t}$ and $j \in \frac{1}{2}\mathbb{Z}$ is the largest index for which $\mathfrak{s}_j = \mathfrak{t}_j$.

If $(\lambda,\ell) \in \hat{A}_i$, let $\mathfrak{t}^\lambda$ denote the maximal element in $\hat{A}^{(\lambda,\ell)}_i$ under the partial order $\succcurlyeq$. Similarly, if $(\mu,m) \in \hat{A}_{i+\frac{1}{2}}$, let $\mathfrak{t}^\mu$ denote the maximal element in $\hat{A}^{(\mu,m)}_{i+\frac{1}{2}}$ under the partial order $\succcurlyeq$. We obtain the next statement by induction on k.

Proposition 3.7. *If* $(\lambda,\ell) \in A_i$, *then*

$$a^{(i)}_{\mathfrak{t}^\lambda} = w_{i-\ell+1,1} w_{i-\ell+2,2} \cdots w_{i,\ell} \qquad \textit{and} \qquad a^{(i+\frac{1}{2})}_{\mathfrak{t}^\lambda} = w_{i-\ell+1,1} w_{i-\ell+2,2} \cdots w_{i,\ell} p^{(\ell)}_\ell.$$

If $i = 1, 2, \ldots,$ and $(\lambda,\ell) \in \hat{A}_i$, then in light of Proposition 3.7, we define

$$f^{(i)}_\lambda = (w_{i-\ell+1,1} w_{i-\ell+2,2} \cdots w_{i,\ell})^* x^{(i)}_{(\lambda,\ell)} w_{i-\ell+1,1} w_{i-\ell+2,2} \cdots w_{i,\ell} = \left(a^{(i)}_{\mathfrak{t}^\lambda}\right)^* x^{(i)}_{(\lambda,\ell)} a^{(i)}_{\mathfrak{t}^\lambda},$$

and write

$$f^{(i)}_{\mathfrak{t}^\lambda} = \left(f^{(i)}_\lambda + A^{\triangleright(\lambda,\ell)}_i\right) \in \mathcal{A}_i / \mathcal{A}^{\triangleright(\lambda,\ell)}_i$$

and

$$f^{(i+\frac{1}{2})}_{\mathfrak{t}^\lambda} = \left(f^{(i)}_\lambda + A^{\triangleright(\lambda,\ell)}_{i+\frac{1}{2}}\right) \in \mathcal{A}_{i+\frac{1}{2}} / \mathcal{A}^{\triangleright(\lambda,\ell)}_{i+\frac{1}{2}}.$$

Example 3.8. Let $\lambda = (2,1)$ and $\ell = 2$. Then $(\lambda,\ell) \in \hat{A}_{5+\frac{1}{2}}$ and

$$\mathfrak{t}^\lambda = (\emptyset,\emptyset,\emptyset,\emptyset,\emptyset,\emptyset,\ydiagram{1},\ydiagram{1},\ydiagram{2},\ydiagram{2},\ydiagram{2,1},\ydiagram{2,1}).$$

If

$$\mathfrak{t} = (\emptyset,\emptyset,\ydiagram{1},\ydiagram{1},\ydiagram{2},\ydiagram{2},\ydiagram{2,1},\ydiagram{2,1},\ydiagram{2,1,1},\ydiagram{2,1},\ydiagram{2,1,1},\ydiagram{2,1}),$$

then $a^{(5+\frac{1}{2})}_{\mathfrak{t}} = 1$ and $a^{(5+\frac{1}{2})}_{\mathfrak{t}^\lambda} = p_4 w_{4,1} p_5 w_{5,2}$. In the diagram presentation of $\mathcal{A}_{5+\frac{1}{2}}$, we have

$$x^{(5+\frac{1}{2})}_{(\lambda,\ell)} =$$



$$x^{(5+\frac{1}{2})}_{(\lambda,\ell)} a^{(5+\frac{1}{2})}_{\mathfrak{t}^\lambda} =$$



and

$$f^{(5)}_\lambda =$$



If $(\lambda,\ell)\in\hat{A}_{i-\frac{1}{2}}$ and $(\mu,m)\in\hat{A}_i$, and $(\lambda,\ell)\to(\mu,m)$, define

$$p^{(i)}_{\lambda\to\mu}=\begin{cases} w_{m,i}, & \text{if } \mu=\lambda,\\ w_{a_j,i}, & \text{if } \mu=\lambda\cup\{(j,\mu_j)\},\end{cases}\tag{3.7}$$

and

$$\rho^{(i)}_{\lambda\to\mu}=\begin{cases} p_m w_{m,i}, & \text{if } \mu=\lambda,\\ \displaystyle\sum_{r=0}^{\lambda_j} w_{a_j-r,a_j} w_{a_j,i}, & \text{if } \mu=\lambda\cup\{(j,\mu_j)\}.\end{cases}\tag{3.8}$$

where in (3.7) and (3.8) we have written

$$a_j=\ell+\sum_{r=1}^{j}\mu_r.$$

Similarly, if $(\lambda,\ell)\in\hat{A}_i$ and $(\mu,m)\in\hat{A}_{i+\frac{1}{2}}$, and $(\lambda,\ell)\to(\mu,m)$, let

$$p^{(i+\frac{1}{2})}_{\lambda\to\mu}=\begin{cases} 1, & \text{if } \lambda=\mu,\\ w_{m,i}p_{i+\frac{1}{2}}w_{i,a_j}\displaystyle\sum_{r=0}^{\mu_j} w_{a_j,a_j-r}, & \text{if } \lambda=\mu\cup\{(j,\lambda_j)\},\end{cases}\tag{3.9}$$

and

$$\rho^{(i+\frac{1}{2})}_{\lambda\to\mu}=\begin{cases} 1, & \text{if } \lambda=\mu,\\ p_m w_{m,i}p_{i+\frac{1}{2}}w_{i,a_j}, & \text{if } \lambda=\mu\cup\{(j,\lambda_j)\},\end{cases}\tag{3.10}$$

where in (3.9) and (3.10) we have written

$$a_j=\ell+\sum_{r=1}^{j}\lambda_r.$$

The next statement is verified using the braid relations.

Proposition 3.9. *(1) If* $(\lambda,\ell)\in\hat{A}_{i-\frac{1}{2}}$, $(\mu,m)\in\hat{A}_i$ *and* $(\lambda,\ell)\to(\mu,m)$, *then*

$$f^{(i)}_{\mu}p^{(i)}_{\lambda\to\mu}=\rho^{(i)}_{\lambda\to\mu}f^{(i-1)}_{\lambda}.$$

(2) If $(\lambda,\ell)\in\hat{A}_i$, $(\mu,m)\in\hat{A}_{i+\frac{1}{2}}$ *and* $(\lambda,\ell)\to(\mu,m)$, *then*

$$f^{(i)}_{\mu}p^{(i+\frac{1}{2})}_{\lambda\to\mu}=\rho^{(i+\frac{1}{2})}_{\lambda\to\mu}f^{(i)}_{\lambda}.$$

For $\mathfrak{t}=(\lambda^{(0)},\lambda^{(\frac{1}{2})},\dots,\lambda^{(k)})\in\hat{A}_k^{(\lambda,\ell)}$, let

$$p_{\mathfrak{t}}=p^{(k)}_{\lambda^{(k-\frac{1}{2})}\to\lambda^{(k)}}p^{(k-\frac{1}{2})}_{\lambda^{(k-1)}\to\lambda^{(k-\frac{1}{2})}}\cdots p^{(\frac{1}{2})}_{\lambda^{(0)}\to\lambda^{(\frac{1}{2})}}.$$

Similarly, for $\mathfrak{s}=(\lambda^{(0)},\lambda^{(\frac{1}{2})},\dots,\lambda^{(k+\frac{1}{2})})\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$, let

$$p_{\mathfrak{s}}=p^{(k+\frac{1}{2})}_{\lambda^{(k)}\to\lambda^{(k+\frac{1}{2})}}p^{(k)}_{\lambda^{(k-\frac{1}{2})}\to\lambda^{(k)}}\cdots p^{(\frac{1}{2})}_{\lambda^{(0)}\to\lambda^{(\frac{1}{2})}}.$$

Let $i\in\mathbb{Z}_{\geqslant 0}$ and $(\lambda,\ell)\in\hat{A}_i$. By Theorems 3.2 and 3.3, we may write

$$A_i^{(\lambda,\ell)}=\{f^{(i)}_{\mathfrak{t}^\lambda}p\mid p\in\mathcal{A}_i\}\qquad\text{and}\qquad A_{i+\frac{1}{2}}^{(\lambda,\ell)}=\{f^{(i)}_{\mathfrak{t}^\lambda}p\mid p\in\mathcal{A}_{i+\frac{1}{2}}\}.$$

Proposition 3.10. *Let* $(\mu,m)\in\hat{A}_i$.

(1) The set $\{f^{(i)}_{\mathfrak{t}^\mu}p_{\mathfrak{t}}\mid\mathfrak{t}\in\hat{A}_i^{(\mu,m)}\}$ *is an* R*–basis for* $A_i^{(\mu,m)}$.

(2) The set $\{f_{\mathfrak{t}^\mu}^{(i+\frac{1}{2})} p_\mathfrak{t} \mid \mathfrak{t} \in \hat{A}^{(\mu,m)}_{i+\frac{1}{2}}\}$ *is an* R*–basis for* $A^{(\mu,m)}_{i+\frac{1}{2}}$.

Proof. (1) If $(\lambda,\ell) \in \hat{A}_{i-\frac{1}{2}}$ and $(\lambda,\ell) \to (\mu,m)$, then

$$f_{\mathfrak{t}^\mu}^{(i)} p^{(i)}_{\lambda\to\mu} = \rho^{(i)}_{\lambda\to\mu} f^{(i-1)}_\lambda + A_i^{\rhd(\mu,m)}. \tag{3.11}$$

If $p \in \mathcal{A}_{i-\frac{1}{2}}$ and $f_\lambda^{(i-\frac{1}{2})} p \in \mathcal{A}^{\rhd(\lambda,\ell)}_{i-\frac{1}{2}}$, then by (1) of Corollary 3.5, there exist $r_\mathfrak{s} \in R$, for $\mathfrak{s} \in \hat{A}_i^{(\mu,m)}$, such that

$$\rho^{(i)}_{\lambda\to\mu} f^{(i-1)}_\lambda p \equiv \sum_{\mathfrak{s}\in \hat{A}_i^{(\mu,m)}} r_\mathfrak{s} f^{(i)}_\mu p^{(i)}_\mathfrak{s} \mod \mathcal{A}_i^{\rhd(\mu,m)}, \tag{3.12}$$

where $r_\mathfrak{s} = 0$ unless $\operatorname{Shape}(\mathfrak{s}|_{i-\frac{1}{2}}) \rhd (\lambda,\ell)$. Let $\{(\lambda^{(j)},\ell_j) \mid j = 1,\dots,t\}$ be an indexing of the set $\{(\lambda,\ell) \in \hat{A}_{i-\frac{1}{2}} \mid (\lambda,\ell) \to (\mu,m)\}$ such that $(\lambda^{(r)},\ell_r) \rhd (\lambda^{(s)},\ell_s)$ whenever $s > r$, and define

$$N_j = \sum_{r=1}^{j} f^{(i)}_{\mathfrak{t}^\mu} p^{(i)}_{\lambda^{(r)}\to\mu} \mathcal{A}_{i-\frac{1}{2}} \qquad\qquad (\text{for } j = 1,\dots,t).$$

By (3.11) and (3.12),

$$\{0\} = N_0 \subseteq N_1 \subseteq \cdots \subseteq N_t = A_i^{(\mu,m)}$$

is a filtration by $\mathcal{A}_{i-\frac{1}{2}}$–modules, and, for $j = 1,\dots,t$, the map

$$\begin{aligned} A^{(\lambda^{(j)},\ell_j)}_{i-\frac{1}{2}} &\longrightarrow N_j/N_{j-1} \\ f^{(i-\frac{1}{2})}_{\mathfrak{t}^{\lambda^{(j)}}} a &\mapsto f^{(i)}_{\mathfrak{t}^\mu} p^{(i)}_{\lambda^{(j)}\to\mu} a + N_{j-1} \qquad (\text{for } a \in \mathcal{A}_{i-\frac{1}{2}}) \end{aligned} \tag{3.13}$$

is an isomorphism of $\mathcal{A}_{i-\frac{1}{2}}$–modules. By induction we may take, for each $(\lambda,\ell) \in \hat{A}_{i-\frac{1}{2}}$, an R–basis for $A^{(\lambda,\ell)}_{i-\frac{1}{2}}$ of the form $\{f^{(i-\frac{1}{2})}_{\mathfrak{t}^\lambda} p_\mathfrak{s} \mid \mathfrak{s} \in \hat{A}^{(\mu,m)}_{i-\frac{1}{2}}\}$. Then, using the map (3.13) to push bases for $A^{(\lambda^{(j)},\ell_j)}_{i-\frac{1}{2}}$ onto bases for N_j/N_{j-1}, completes the proof of (1).

(2) Regard (μ,m) as an element of $\hat{A}_{i+\frac{1}{2}}$. If $(\lambda,\ell) \in \hat{A}_i$ and $(\lambda,\ell) \to (\mu,m)$, then

$$f_{\mathfrak{t}^\mu}^{(i+\frac{1}{2})} p^{(i+\frac{1}{2})}_{\lambda\to\mu} = \rho^{(i+\frac{1}{2})}_{\lambda\to\mu} f^{(i)}_\lambda + A_{i+\frac{1}{2}}^{\rhd(\mu,m)}. \tag{3.14}$$

If $p \in \mathcal{A}_i$ and $f^{(i)}_\lambda p \in \mathcal{A}_i^{\rhd(\lambda,\ell)}$, then by item (2) of Corollary 3.5, there exist $r_\mathfrak{s} \in R$, for $\mathfrak{s} \in \hat{A}^{(\mu,m)}_{i+\frac{1}{2}}$, such that

$$\rho^{(i+\frac{1}{2})}_{\lambda\to\mu} f^{(i)}_\lambda p \equiv \sum_{\mathfrak{s}\in \hat{A}^{(\mu,m)}_{i+\frac{1}{2}}} r_\mathfrak{s} f^{(i)}_\mu p^{(i+\frac{1}{2})}_\mathfrak{s} \mod \mathcal{A}^{\rhd(\mu,m)}_{i+\frac{1}{2}} \tag{3.15}$$

where $r_\mathfrak{s} = 0$ unless $\operatorname{Shape}(\mathfrak{s}|_i) \rhd (\lambda,\ell)$. Let $\{(\lambda^{(j)},\ell_j) \mid j = 1,\dots,t\}$ be an indexing of the set $\{(\lambda,\ell) \in \hat{A}_i \mid (\lambda,\ell) \to (\mu,m)\}$ such that $(\lambda^{(r)},\ell_r) \rhd (\lambda^{(s)},\ell_s)$ whenever $s > r$, and define

$$N_j = \sum_{r=1}^{j} f^{(i+\frac{1}{2})}_{\mathfrak{t}^\mu} p_{\lambda^{(r)}\to\mu} \mathcal{A}_i \qquad\qquad (\text{for } j = 1,\dots,t).$$

By (3.14) and (3.15),

$$\{0\} = N_0 \subseteq N_1 \subseteq \cdots \subseteq N_t = A^{(\mu,m)}_{i+\frac{1}{2}}$$

is a filtration by $\mathcal{A}_i$–modules, and, for $j = 1, \ldots, t$, the map

$$\begin{aligned} A_i^{(\lambda_j,\ell_j)} &\longrightarrow N_j/N_{j-1} \\ f^{(i)}_{\mathfrak{t}^{\lambda^{(j)}}}\, a &\mapsto f^{(i)}_{\mathfrak{t}^{\mu}} p^{(i+\frac{1}{2})}_{\lambda^{(j)}\to\mu} a + N_{j-1} \qquad (\text{for } a \in \mathcal{A}_i) \end{aligned} \tag{3.16}$$

is an isomorphism of $\mathcal{A}_i$–modules. Using induction as in (1) above completes the proof of (2). $\square$

In Theorems 3.11 and 3.12 the bases (3.3) and (3.4) are expressed relative to the elements $f^{(i)}_{\lambda}$, for $(\lambda, \ell) \in \hat{\mathcal{A}}_i$.

Theorem 3.11. *If $i = 1, 2, \ldots,$ then the set*

$$\mathscr{A}_i = \{p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{\mathcal{A}}_i^{(\lambda,\ell)} \text{ and } (\lambda, \ell) \in \hat{\mathcal{A}}_i\}$$

is an R–basis for $\mathcal{A}_i(z)$. Moreover the following statements hold:

(1) If $(\lambda, \ell) \in \hat{\mathcal{A}}_i$, $\mathfrak{t} \in \hat{\mathcal{A}}_k^{(\lambda,\ell)}$ and $p \in \mathcal{A}_k(z)$, then there exist $r_{\mathfrak{v}} \in R$, for $\mathfrak{v} \in \hat{\mathcal{A}}_i^{(\lambda,\ell)}$, such that

$$p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{t}} p \equiv \sum_{\mathfrak{v} \in \hat{A}_k^{(\lambda,\ell)}} r_{\mathfrak{v}} p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{v}} \mod \mathcal{A}_i^{\rhd(\lambda,\ell)} \qquad \text{for all } \mathfrak{s} \in \hat{A}_i^{(\lambda,\ell)}, \tag{3.17}$$

where $\mathcal{A}_i^{\rhd(\lambda,\ell)}$, for $(\lambda, \ell) \in \hat{\mathcal{A}}_i$, is the R–module freely generated by

$$\big\{p^*_{\mathfrak{s}} f^{(i)}_{\mu} p_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{A}_i^{(\mu,m)} \text{ for } (\mu, m) \in \hat{A}_i \text{ and } (\mu, m) \rhd (\lambda, \ell)\big\}.$$

(2) If $(\lambda, \ell) \in \hat{\mathcal{A}}_i$ and $\mathfrak{s}, \mathfrak{t} \in \hat{A}_i^{(\lambda,\ell)}$, then $* : p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{t}} \mapsto p^*_{\mathfrak{t}} f^{(i)}_{\lambda} p_{\mathfrak{s}}$.

Theorem 3.12. *If $i = 1, 2, \ldots,$ then the set*

$$\mathscr{A}_{i+\frac{1}{2}} = \{p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{\mathcal{A}}_{i+\frac{1}{2}}^{(\lambda,\ell)} \text{ and } (\lambda, \ell) \in \hat{\mathcal{A}}_{i+\frac{1}{2}}\}$$

is an R–basis for $\mathcal{A}_{i+\frac{1}{2}}(z)$. Moreover the following statements hold:

(1) If $(\lambda, \ell) \in \hat{\mathcal{A}}_{i+\frac{1}{2}}$, $\mathfrak{t} \in \hat{\mathcal{A}}_{i+\frac{1}{2}}^{(\lambda,\ell)}$ and $p \in \mathcal{A}_{i+\frac{1}{2}}(z)$, then there exist $r_{\mathfrak{v}} \in R$, for $\mathfrak{v} \in \hat{\mathcal{A}}_{i+\frac{1}{2}}^{(\lambda,\ell)}$, such that

$$p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{t}} p \equiv \sum_{\mathfrak{v} \in \hat{A}_{i+\frac{1}{2}}^{(\lambda,\ell)}} r_{\mathfrak{v}} p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{v}} \mod \mathcal{A}_{i+\frac{1}{2}}^{\rhd(\lambda,\ell)} \qquad \text{for all } \mathfrak{s} \in \hat{A}_{i+\frac{1}{2}}^{(\lambda,\ell)}, \tag{3.18}$$

where $\mathcal{A}_{i+\frac{1}{2}}^{\rhd(\lambda,\ell)}$, for $(\lambda, \ell) \in \hat{\mathcal{A}}_{i+\frac{1}{2}}$, is the R–module freely generated by

$$\big\{p^*_{\mathfrak{s}} f^{(i)}_{\mu} p_{\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \hat{A}_{i+\frac{1}{2}}^{(\mu,m)} \text{ for } (\mu, m) \in \hat{A}_{i+\frac{1}{2}} \text{ and } (\mu, m) \rhd (\lambda, \ell)\big\}.$$

(2) If $(\lambda, \ell) \in \hat{\mathcal{A}}_{i+\frac{1}{2}}$ and $\mathfrak{s}, \mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$, then $* : p^*_{\mathfrak{s}} f^{(i)}_{\lambda} p_{\mathfrak{t}} \mapsto p^*_{\mathfrak{t}} f^{(i)}_{\lambda} p_{\mathfrak{s}}$.

If $a = (i, j)$ is a node, let $c(a) = j - i$ be the content of a. Let $(\mu, m) \in \hat{A}_k$. For $\mathfrak{t} = (\mu^{(0)}, \mu^{(\frac{1}{2})}, \ldots, \mu^{(k)}) \in \hat{A}_k^{(\mu,m)}$, and $i = 1, \ldots, k$, define

$$c_{\mathfrak{t}}(i - \tfrac{1}{2}) = \begin{cases} |\mu^{(i-1)}|, & \text{if } \mu^{(i-1)} = \mu^{(i-\frac{1}{2})}; \\ z - c(a), & \text{if } \mu^{(i-\frac{1}{2})} = \mu^{(i-1)} \setminus \{a\}. \end{cases}$$

$$c_{\mathfrak{t}}(i) = \begin{cases} z - |\mu^{(i)}|, & \text{if } \mu^{(i)} = \mu^{(i-\frac{1}{2})}, \\ c(a), & \text{if } \mu^{(i)} = \mu^{(i-\frac{1}{2})} \cup \{a\}, \text{ and} \end{cases}$$

If $\mathfrak{s} = (\mu^{(0)}, \mu^{(\frac{1}{2})}, \dots, \mu^{(k+\frac{1}{2})}) \in \hat{A}^{(\mu,m)}_{k+\frac{1}{2}}$ and $\mathfrak{s}|_k = \mathfrak{t}$, let

$$c_{\mathfrak{t}}(k+\tfrac{1}{2}) = \begin{cases} |\mu^{(k)}|, & \text{if } \mu^{(k)} = \mu^{(k+\frac{1}{2})}; \\ z - c(a), & \text{if } \mu^{(k+\frac{1}{2})} = \mu^{(k)} \setminus \{a\}, \end{cases}$$

and let $c_{\mathfrak{s}}(i-\frac{1}{2}) = c_{\mathfrak{t}}(i-\frac{1}{2})$ and $c_{\mathfrak{s}}(i) = c_{\mathfrak{t}}(i)$ for $i = 1, \dots, k$.

The next statement shows that the elements $\{L_i, L_{i+\frac{1}{2}} \mid i = 0, 1, \dots\}$ are a additive Jucys–Murphy family in the sense of Goodman and Graber (Definition 3.4 of [GG1]).

Proposition 3.13. *For $(\lambda,\ell) \in \hat{A}_k$, the following statements hold.*

(1) If $i = 1, 2, \dots, k$, then

$$f^{(k)}_\lambda L_{i-\frac{1}{2}} \equiv c_{\mathfrak{t}^\lambda}(i-\tfrac{1}{2}) f^{(k)}_\lambda \mod \mathcal{A}_k^{\rhd(\lambda,\ell)} \quad \text{and} \quad f^{(k)}_\lambda L_i \equiv c_{\mathfrak{t}^\lambda}(i) f^{(k)}_\lambda \mod \mathcal{A}_k^{\rhd(\lambda,\ell)} \tag{3.19}$$

and

$$f^{(k)}_\lambda L_i \equiv c_{\mathfrak{t}^\lambda}(i) f^{(k)}_\lambda \mod \mathcal{A}_{k+\frac{1}{2}}^{\rhd(\lambda,\ell)} \quad \text{and} \quad f^{(k)}_\lambda L_{i+\frac{1}{2}} \equiv c_{\mathfrak{t}^\lambda}(i+\tfrac{1}{2}) f^{(k)}_\lambda \mod \mathcal{A}_{k+\frac{1}{2}}^{\rhd(\lambda,\ell)}. \tag{3.20}$$

(2) The central element $z_k = L_1 + L_{1+\frac{1}{2}} + L_2 + \dots + L_k \in \mathcal{A}_k$ acts on $A_k^{(\lambda,\ell)}$ as a scalar multiple of the identity by

$$\ell z + \sum_{b\in\lambda} c(b) + \frac{(k-\ell)(k-\ell-1)}{2}.$$

(3) The central element $z_{k+\frac{1}{2}} = L_1 + L_{1+\frac{1}{2}} + L_2 + \dots + L_{k+\frac{1}{2}} \in \mathcal{A}_{k+\frac{1}{2}}$ acts on $A_{k+\frac{1}{2}}^{(\lambda,\ell)}$ as a scalar multiple of the identity by

$$\ell z + \sum_{b\in\lambda} c(b) + \frac{(k-\ell+1)(k-\ell)}{2}.$$

Proof. We prove (1). For $j = 1, 2, \dots$, define $\{L_i^{(j)} \mid j = 0, 1, \dots\}$ and $\{L_{i+\frac{1}{2}}^{(j)} \mid i = 0, 1, \dots\}$ by

$$L^{(1)}_{i-\frac{1}{2}} = L_{i-\frac{1}{2}}, \qquad \text{and} \qquad L^{(j+1)}_{i-\frac{1}{2}} = w_{j,j+i} L^{(j)}_{i-\frac{1}{2}} w_{j+i,j}, \qquad \text{for } j = 1, 2, \dots,$$

and

$$L^{(1)}_{i} = L_{i}, \qquad \text{and} \qquad L^{(j+1)}_{i} = w_{j,j+i} L^{(j)}_{i} w_{j+i,j}, \qquad \text{for } j = 1, 2, \dots.$$

Since $p_1 L_1 = z p_1$ and $p_1 L_{1+\frac{1}{2}} = 0$, using induction, we obtain

$$p_j L_i^{(j)} = \begin{cases} 0, & \text{if } i = 0, \\ z p_j, & \text{if } i = 1, \\ p_j L_{i-1}^{(j+1)}, & \text{if } i \geqslant 2, \end{cases} \qquad \text{and} \qquad p_j L_{i+\frac{1}{2}}^{(j)} = \begin{cases} 0, & \text{if } i = 0, \\ 0, & \text{if } i = 1, \\ p_j L_{i-\frac{1}{2}}^{(j+1)}, & \text{if } i \geqslant 2, \end{cases}$$

so that

$$p_\ell^{(\ell)} L_i^{(j)} = \begin{cases} 0, & \text{if } i = 0, \\ z p_\ell^{(\ell)}, & \text{if } 1 \leqslant i \leqslant \ell, \\ p_\ell^{(\ell)} L_{i-\ell}^{(\ell+1)}, & \text{if } \ell < i, \end{cases} \qquad \text{and} \qquad p_\ell^{(\ell)} L_{i+\frac{1}{2}}^{(j)} = \begin{cases} 0, & \text{if } i = 0, \\ 0, & \text{if } 1 \leqslant i \leqslant \ell, \\ p_\ell^{(\ell)} L_{i-\ell+\frac{1}{2}}^{(\ell+1)}, & \text{if } \ell < i. \end{cases}$$

Thus

$$f^{(k)}_\lambda L_i = c_{\mathfrak{t}^\lambda}(i) f^{(k)}_\lambda \qquad \text{and} \qquad f^{(k)}_\lambda L_{i+\frac{1}{2}} = c_{\mathfrak{t}^\lambda}(i+\tfrac{1}{2}) f^{(k)}_\lambda \qquad \text{for } i = 1, \dots, \ell,$$

as required. Now suppose that $\ell < i \leqslant k$. Since the image of $L_{i-\ell}$ under the quotient map $A_{k-\ell} \to R\mathfrak{S}_{k-\ell}$ is the $(i-\ell)$–th Jucys–Murphy element in $\mathfrak{S}_{k-\ell}$, we have

$$c_\lambda L_{i-\ell} \equiv c_{\mathfrak{t}^\lambda}(i) c_\lambda \mod A_{k-\ell}^{\rhd(\lambda,0)}.$$

As $p_k^{(\ell)} A_{k-\ell}^{\rhd(\lambda,0)} \subseteq A_k^{\rhd(\lambda,\ell)}$, it now follows that

$$f_\lambda^{(k)} L_i = f_\lambda^{(k)} L_{i-\ell}^{(\ell+1)} \equiv c_{\mathfrak{t}^\lambda}(i) f_\lambda^{(k)} \mod A_k^{\rhd(\lambda,\ell)},$$

as required. If $\ell < i \leqslant k$, then the image of $L_{i-\ell+\frac{1}{2}}$ under the quotient map $A_{k-\ell+\frac{1}{2}} \to R\mathfrak{S}_{k-\ell}$ is the identity in $\mathfrak{S}_{k-\ell}$, and the above argument also shows that $f_\lambda^{(k)} L_{i+\frac{1}{2}} \equiv f_\lambda^{(k)} \mod A_k^{\rhd(\lambda,\ell)}$. This verifies (3.19). The proof of (3.20) is identical.

Writing

$$f_\lambda^{(k)} z_k = \sum_{i=1}^{\ell} f_\lambda^{(k)} (L_{i-\frac{1}{2}} + L_i) + \sum_{i=\ell+1}^{k} f_\lambda^{(k)} (L_{i-\frac{1}{2}} + L_i),$$

and using (1), proves (2) and (3). □

The next statement which shows that the family $\{L_i, L_{i+\frac{1}{2}}, | \ i = 1, \dots, k\}$ acts triangularly on $\mathcal{A}_{k+\frac{1}{2}}$, now follows by a direct calculation or by Theorem 3.7 of [GG1].

Corollary 3.14. *If $(\lambda,\ell) \in \hat{A}_k$ and $\mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ then the following statements hold.*

(1) If $i = 1, \dots, k$, then there exist $r_\mathfrak{u} \in R$, for $\mathfrak{u} \in \hat{A}_k^{(\lambda,\ell)}$ with $\mathfrak{u} \succ \mathfrak{t}$, such that

$$f_{\mathfrak{t}^\lambda}^{(i+\frac{1}{2})} p_\mathfrak{t} L_i = c_\mathfrak{t}(i) f_{\mathfrak{t}^\lambda}^{(i)} p_\mathfrak{t} + \sum_{\mathfrak{u} \succ \mathfrak{t}} r_\mathfrak{u} f_{\mathfrak{t}^\lambda}^{(i+\frac{1}{2})} p_\mathfrak{u}$$

(2) If $i = 1, \dots, k$, then there exist $r_\mathfrak{u} \in R$, for $\mathfrak{u} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ with $\mathfrak{u} \succ \mathfrak{t}$, such that

$$f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_\mathfrak{t} L_{i+\frac{1}{2}} = c_\mathfrak{t}(k+\tfrac{1}{2}) f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_\mathfrak{t} + \sum_{\mathfrak{u} \succ \mathfrak{t}} r_\mathfrak{u} f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_\mathfrak{u}.$$

Using the $\mathcal{A}_k$–module embedding $A_k^{(\lambda,\ell)} \subseteq A_{k+\frac{1}{2}}^{(\lambda,\ell)}$, for $(\lambda,\ell) \in \hat{A}_k$, Corollary 3.14 shows that the family $\{L_{i-\frac{1}{2}}, L_i \mid i = 1, \dots, k\}$ acts triangularly on $\mathcal{A}_k$.

4. A Seminormal Form

In this section, we use the triangular action of the Jucys–Murphy elements $\{L_i, L_{i+\frac{1}{2}} \mid i = 1, \dots, k\}$ to define a seminormal form for the partition algebras. Let κ denote the field of fractions of $R = \mathbb{Z}[z]$, and

$$A_k(z) = \mathcal{A}_k(z) \otimes_R \kappa \qquad \text{and} \qquad A_{k+\frac{1}{2}}(z) = \mathcal{A}_{k+\frac{1}{2}}(z) \otimes_R \kappa$$

denote partition algebras over κ. We write $A_k = A_k(z)$ and $A_{k+\frac{1}{2}} = A_{k+\frac{1}{2}}(z)$. If $(\lambda,\ell) \in \hat{A}_k$, let

$$A_k^{\rhd(\lambda,\ell)} = \mathcal{A}_k^{\rhd(\lambda,\ell)} \otimes_R \kappa \qquad \text{and} \qquad A_{k+\frac{1}{2}}^{\rhd(\lambda,\ell)} = \mathcal{A}_{k+\frac{1}{2}}^{\rhd(\lambda,\ell)} \otimes_R \kappa.$$

If $(\lambda,\ell) \in \hat{A}_k$, there will be no confusion in letting $A_k^{(\lambda,\ell)}$ and $A_{k+\frac{1}{2}}^{(\lambda,\ell)}$ denote the cell modules for A_k and $A_{k+\frac{1}{2}}$ respectively.

If $(\lambda,\ell)\in\hat{A}_{k+\frac12}$, and $\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac12}$, define

$$F_{\mathfrak{t}}=\prod_{i\in\{\frac12,1,\ldots,k+\frac12\}}\ \prod_{\substack{\mathfrak{u}\in\hat{A}^{(\lambda,\ell)}_{k+\frac12}\\ c_{\mathfrak{u}}(i)\neq c_{\mathfrak{t}}(i)}}\frac{L_i-c_{\mathfrak{u}}(i)}{c_{\mathfrak{t}}(i)-c_{\mathfrak{u}}(i)},$$

and

$$f^{(k+\frac12)}_{\mathfrak{t}}=f^{(k)}_{\mathfrak{t}^\lambda}p_{\mathfrak{t}}F_{\mathfrak{t}},\qquad\text{and}\qquad F_{\mathfrak{s}\mathfrak{t}}=F_{\mathfrak{s}}p^*_{\mathfrak{s}}f^{(k)}_{\lambda}p_{\mathfrak{t}}F_{\mathfrak{t}}.$$

Following §3 of [Mat], or by direct calculation, we obtain the next statement.

Proposition 4.1. *Let $k=1,2,\ldots,$ and $(\mu,m)\in\hat{A}_{k+\frac12}$.*

(1) If $\mathfrak{t}\in\hat{A}^{(\mu,m)}_{k+\frac12}$, then there exist $r_{\mathfrak{s}}\in\kappa$, for $\mathfrak{s}\in\hat{A}^{(\mu,m)}_{k+\frac12}$, such that

$$f^{(k+\frac12)}_{\mathfrak{t}}=f^{(k+\frac12)}_{\mathfrak{t}^\mu}p_{\mathfrak{t}}+\sum_{\mathfrak{s}\in\hat{A}^{(\mu,m)}_{k+\frac12}}r_{\mathfrak{s}}f^{(k+\frac12)}_{\mathfrak{t}^\mu}p_{\mathfrak{s}},\qquad\text{where } r_{\mathfrak{s}}=0 \text{ unless } \mathfrak{s}\succ\mathfrak{t}.$$

(2) The sets

$$\big\{f^{(k+\frac12)}_{\mathfrak{t}}\mid\mathfrak{t}\in\hat{A}^{(\mu,m)}_{k+\frac12}\big\}\qquad\text{and}\qquad\big\{F_{\mathfrak{s}\mathfrak{t}}\mid\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\mu,m)}_{k+\frac12}\text{ and }(\mu,m)\in\hat{A}_{k+\frac12}\big\}$$

are bases over κ for $A^{(\mu,m)}_{k+\frac12}$ and $A_{k+\frac12}$ respectively.

(3) If $\mathfrak{t}\in\hat{A}^{(\mu,m)}_{k+\frac12}$, then

$$f^{(k+\frac12)}_{\mathfrak{t}}L_i=c_{\mathfrak{t}}(i)f^{(k+\frac12)}_{\mathfrak{t}}\qquad\text{and}\qquad f^{(k+\frac12)}_{\mathfrak{t}}L_{i+\frac12}=c_{\mathfrak{t}}(i+\tfrac12)f^{(k+\frac12)}_{\mathfrak{t}}\qquad\text{for } i=0,1,\ldots,k.$$

(4) If $(\lambda,\ell)\in\hat{A}_{k+\frac12}$ then the symmetric contravariant bilinear form $\langle\,,\rangle:A^{(\mu,m)}_{k+\frac12}\times A^{(\lambda,\ell)}_{k+\frac12}\to\kappa$, defined by

$$\langle f^{(k+\frac12)}_{\mathfrak{t}^\mu}p_{\mathfrak{s}},f^{(k+\frac12)}_{\mathfrak{t}^\mu}p_{\mathfrak{t}}\rangle f^{(k)}_{\mu}\equiv f^{(k)}_{\mu}p_{\mathfrak{s}}p^*_{\mathfrak{t}}f^{(k)}_{\mu}\mod A^{\rhd(\mu,m)}_{k+\frac12},\qquad\text{for }\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\mu,m)}_{k+\frac12},\tag{4.1}$$

satisfies

$$\langle f^{(k+\frac12)}_{\mathfrak{s}},f^{(k+\frac12)}_{\mathfrak{t}}\rangle=\delta_{\mathfrak{s}\mathfrak{t}}\langle f^{(k+\frac12)}_{\mathfrak{t}},f^{(k+\frac12)}_{\mathfrak{t}}\rangle,\qquad\text{for }\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\mu,m)}_{k+\frac12}.$$

(5) If $\mathfrak{s},\mathfrak{t},\mathfrak{u}\in\hat{A}^{(\mu,m)}_{k+\frac12}$, then

$$f^{(k+\frac12)}_{\mathfrak{s}}F_{\mathfrak{t}}=\delta_{\mathfrak{s}\mathfrak{t}}f^{(k+\frac12)}_{\mathfrak{s}}\qquad\text{and}\qquad f_{\mathfrak{u}}F_{\mathfrak{s}\mathfrak{t}}=\langle f^{(k+\frac12)}_{\mathfrak{s}},f^{(k+\frac12)}_{\mathfrak{u}}\rangle f^{(k+\frac12)}_{\mathfrak{t}}.$$

(6) If $\mathfrak{t},\mathfrak{u}\in\hat{A}^{(\mu,m)}_{k+\frac12}$, then

$$F_{\mathfrak{t}}F_{\mathfrak{u}}=\delta_{\mathfrak{t}\mathfrak{u}}F_{\mathfrak{t}}\qquad\text{and}\qquad F^2_{\mathfrak{s}\mathfrak{t}}=\langle f^{(k+\frac12)}_{\mathfrak{s}},f^{(k+\frac12)}_{\mathfrak{t}}\rangle F_{\mathfrak{s}\mathfrak{t}}.$$

(7) If $(\mu,m)\in\hat{A}_{k+\frac12}$, then there is an A_k–module isomorphism

$$A^{(\mu,m)}_{k+\frac12}\cong\bigoplus_{\substack{(\lambda,\ell)\in\hat{A}_k\\(\lambda,\ell)\to(\mu,m)}}A^{(\lambda,\ell)}_{k},$$

determined by the maps (3.16).

If $(\mu, m) \in \hat{A}_k$, the homomorphism (3.16) determines an inclusion $A_k^{(\mu,m)} \hookrightarrow A_{k+\frac{1}{2}}^{(\mu,m)}$ of $\mathcal{A}_k$–modules. Therefore, in what follows, we identify the seminormal form of the A_k–module $A_k^{(\mu,m)}$, for $(\mu, m) \in \hat{A}_k$, with the κ–vector space generated by

$$\left\{ f_{\mathfrak{s}}^{(k)} = f_{\mathfrak{t}}^{(k+\frac{1}{2})} \;\middle|\; \mathfrak{s} \in \hat{A}_k^{(\mu,m)}, \mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\mu,m)} \text{ and } \mathfrak{s} = \mathfrak{t}|_k \right\},$$

with right A_k–action given by the inclusion $A_k \subseteq A_{k+\frac{1}{2}}$. The $\mathcal{A}_{k-\frac{1}{2}}$–module homomorphisms (3.13) determine an $A_{k-\frac{1}{2}}$–module isomorphism

$$A_k^{(\mu,m)} \cong \bigoplus_{\substack{(\lambda,\ell)\in\hat{A}_{k-\frac{1}{2}} \\ (\lambda,\ell)\to(\mu,m)}} A_{k-\frac{1}{2}}^{(\lambda,\ell)}.$$

If $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$ and $\mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$, define the seminormal matrix entries

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_i = \sum_{\mathfrak{s}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}} p_i(\mathfrak{s},\mathfrak{t}) f_{\mathfrak{s}}^{(k+\frac{1}{2})} \quad \text{and} \quad f_{\mathfrak{t}} p_{i+\frac{1}{2}} = \sum_{\mathfrak{s}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}} p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t}) f_{\mathfrak{s}}^{(k+\frac{1}{2})} \quad \text{for } i = 1,\ldots,k,$$

and

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})} \sigma_{i+\frac{1}{2}} = \sum_{\mathfrak{s}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}} \sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t}) f_{\mathfrak{s}}^{(k+\frac{1}{2})}, \qquad \text{for } i = 1,\ldots,k,$$

and

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})} \sigma_{i+1} = \sum_{\mathfrak{s}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}} \sigma_{i+1}(\mathfrak{s},\mathfrak{t}) f_{\mathfrak{s}}^{(k+\frac{1}{2})}, \qquad \text{for } i = 1,\ldots,k-1.$$

The matrix entries of the transposition s_i in the seminormal representation of $A_{k+\frac{1}{2}}(z)$ are obtained by the relation

$$s_i(\mathfrak{s},\mathfrak{t}) = \sum_{\mathfrak{u}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}} \sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{u}) \sigma_{i+1}(\mathfrak{u},\mathfrak{t}), \qquad \text{for } \mathfrak{s},\mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}, \text{ and } i = 1,\ldots,k-1.$$

In [En2] we have explicitly computed the bilinear form (4.1) with respect to the seminormal basis for $A_{k+\frac{1}{2}}$. We state the results of these calculations below for reference in Theorem 5.1.

Recall that if μ is a partition, then $A(\mu)$ is the set of addable nodes of μ and $R(\mu)$ is the set of removable nodes of μ. If $a = (i, j)$ and $b = (k, \ell)$ are nodes, write $a < b$ if $k < i$, and write $a > b$ if $k > i$. If μ is a partition and a is a node, write

$$A(\mu)^{<a} = \{b \in A(\mu) \mid b < a\} \qquad \text{and} \qquad A(\mu)^{>a} = \{b \in A(\mu) \mid b > a\},$$

and

$$R(\mu)^{<a} = \{b \in R(\mu) \mid b < a\} \qquad \text{and} \qquad R(\mu)^{>a} = \{b \in R(\mu) \mid b > a\}.$$

If $(\lambda, \ell) \in \hat{A}_{i-\frac{1}{2}}$, $(\mu, m) \in \hat{A}_i$, and $(\lambda, \ell) \to (\mu, m)$, define

$$\gamma_{\lambda\to\mu}^{(i)} = \begin{cases} \dfrac{\prod_{b\in A(\mu)}(z - c(b) - |\mu|)}{\prod_{b\in R(\mu)}(z - c(b) - |\mu|)}, & \text{if } \mu = \lambda, \\[2ex] \dfrac{\prod_{b\in A(\mu)^{<a}}(c(a) - c(b))}{\prod_{b\in R(\mu)^{<a}}(c(a) - c(b))}, & \text{if } \mu = \lambda \cup \{a\}. \end{cases}$$

Similarly, if $(\lambda,\ell)\in\hat{A}_i$, $(\mu,m)\in\hat{A}_{i+\frac{1}{2}}$, and $(\lambda,\ell)\to(\mu,m)$, let

$$\gamma^{(i+\frac{1}{2})}_{\lambda\to\mu}=\begin{cases}1, & \text{if } \mu=\lambda,\\ \dfrac{(z-c(a)-|\lambda|)}{(z-c(a)-|\mu|)}\dfrac{(z-c(a)-|\mu|+r)}{(z-c(a)-|\lambda|+r)} & \\ \qquad\times\dfrac{(z+c(a)+|\lambda|-r)}{(z+c(a)+|\mu|-r)}\dfrac{\prod_{b\in R(\mu)^{>a}}(c(a)-c(b))}{\prod_{b\in A(\mu)^{>a}}(c(a)-c(b))}, & \text{if } \lambda=\mu\cup\{a\},\end{cases}$$

where in the last case we have written $a=(j,\lambda_j)$, and

$$r=\sum_{s>j}\mu_s.$$

Theorem 4.2. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$. *If* $\mathfrak{t}=\big(\lambda^{(0)},\lambda^{(\frac{1}{2})},\dots,\lambda^{(k+\frac{1}{2})}\big)\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, *then*

$$\langle f^{(k+\frac{1}{2})}_{\mathfrak{t}},f^{(k+\frac{1}{2})}_{\mathfrak{t}}\rangle=\prod_{i=1}^{k}\gamma^{(i)}_{\lambda^{(i-\frac{1}{2})}\to\lambda^{(i)}}\gamma^{(i+\frac{1}{2})}_{\lambda^{(i)}\to\lambda^{(i+\frac{1}{2})}}. \tag{4.2}$$

5. MAIN RESULTS

Theorems 5.1, 5.2 and 5.3 below give explicit combinatorial formulae for images of generators of the partition algebras in Theorem 2.2 in the seminormal representations. The statement of these results is split into three parts merely to avoid clashes between indices. In Theorem 5.1 we have used the abbreviation $f_{\mathfrak{t}}=f^{(k+\frac{1}{2})}_{\mathfrak{t}}$, for $\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$.

Theorem 5.1. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$. *If* $\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$ *and* $\mathfrak{s}\neq\mathfrak{t}$, *and* $i=1,2,\dots,k$, *then*

$$p_i(\mathfrak{s},\mathfrak{t})=\sqrt{\frac{p_i(\mathfrak{s},\mathfrak{s})p_i(\mathfrak{t},\mathfrak{t})\langle f_{\mathfrak{t}},f_{\mathfrak{t}}\rangle}{\langle f_{\mathfrak{s}},f_{\mathfrak{s}}\rangle}}\quad\text{and}\quad p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})=\sqrt{\frac{p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{s})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})\langle f_{\mathfrak{t}},f_{\mathfrak{t}}\rangle}{\langle f_{\mathfrak{s}},f_{\mathfrak{s}}\rangle}}, \tag{5.1}$$

where

$$p_i(\mathfrak{t},\mathfrak{t})=\begin{cases}\dfrac{\prod_{a\in A(\mu)}\big(z-c(a)-|\mu|\big)}{\prod_{b\in R(\mu)}\big(z-c(b)-|\mu|\big)}, & \text{if } \mathfrak{t}^{(i-1)}=\mathfrak{t}^{(i-\frac{1}{2})}=\mathfrak{t}^{(i)}=\mu,\\ \dfrac{(z-c(\beta)-|\mu|+1)}{(z-c(\beta)-|\mu|)}\dfrac{\prod_{a\in A(\mu)}(c(\beta)-c(a))}{\prod_{\substack{b\in R(\mu),\\ b\neq\beta}}(c(\beta)-c(b))}, & \text{if } \mathfrak{t}^{(i-1)}=\mathfrak{t}^{(i)}=\mu \text{ and } \mathfrak{t}^{(i-\frac{1}{2})}=\mu\setminus\{\beta\},\\ 0, & \text{otherwise},\end{cases}$$

and,

$$p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})=\begin{cases}\dfrac{\prod_{b\in R(\mu)}(z-c(b)-|\mu|)}{\prod_{a\in A(\mu)}(z-c(a)-|\mu|)}, & \text{if } \mathfrak{t}^{(i-\frac{1}{2})}=\mathfrak{t}^{(i)}=\mathfrak{t}^{(i+\frac{1}{2})}=\mu,\\ \dfrac{(z+c(\beta)+|\mu|+1)}{(z+c(\beta)+|\mu|)}\dfrac{\prod_{b\in R(\mu)}(c(\beta)-c(b))}{\prod_{\substack{a\in A(\mu),\\ a\neq\beta}}(c(\beta)-c(a))}, & \text{if } \mathfrak{t}^{(i-\frac{1}{2})}=\mathfrak{t}^{(i+\frac{1}{2})}=\mu \text{ and } \mathfrak{t}^{(i)}=\mu\cup\{\beta\},\\ 0, & \text{otherwise}.\end{cases}$$

Theorem 5.2. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$ *and* $\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$. *For* $i=1,\dots,k$, *the following statements hold:*

(1) If $\mathfrak{t}^{(i-\frac{1}{2})}=\mathfrak{t}^{(i+\frac{1}{2})}$*, and* $\mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$ *satisfies* $\mathfrak{v}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}$ *and* $\mathfrak{v}^{(i-1)}=\mathfrak{v}^{(i)}$*, then*

$$\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})=\frac{\delta_{\mathfrak{s}\mathfrak{t}}+\big(z-c_{\mathfrak{s}}(i)-c_{\mathfrak{t}}(i-\frac{1}{2})\big)p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})-p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{v})p_i(\mathfrak{v},\mathfrak{t})}{c_{\mathfrak{s}}(i+\frac{1}{2})-c_{\mathfrak{t}}(i-\frac{1}{2})},\qquad \textit{if } \mathfrak{s}\neq\mathfrak{v},$$

and

$$\sigma_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})=\frac{p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})}{p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{v})p_i(\mathfrak{v},\mathfrak{v})}c_{\mathfrak{t}}(i)-\sum_{\substack{\mathfrak{u}\overset{i}{\sim}\mathfrak{v}\\ \mathfrak{u}\neq\mathfrak{v}}}\frac{p_i(\mathfrak{v},\mathfrak{u})\sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t})}{p_i(\mathfrak{v},\mathfrak{v})}.$$

(2) If $\mathfrak{t}^{(i-1)}=\mathfrak{t}^{(i)}$ *and* $\mathfrak{t}^{(i-\frac{1}{2})}\neq\mathfrak{t}^{(i+\frac{1}{2})}$*, and* $\mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$ *satisfies* $\mathfrak{v}\overset{i}{\sim}\mathfrak{t}$ *and* $\mathfrak{v}^{(i-\frac{1}{2})}=\mathfrak{v}^{(i+\frac{1}{2})}$*, then*

$$\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})=\frac{\delta_{\mathfrak{s}\mathfrak{t}}-p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{v})p_i(\mathfrak{v},\mathfrak{t})}{c_{\mathfrak{s}}(i+\frac{1}{2})-c_{\mathfrak{t}}(i-\frac{1}{2})},\qquad \textit{if } \mathfrak{s}\neq\mathfrak{v},$$

and

$$\sigma_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})=-\sum_{\substack{\mathfrak{u}\overset{i}{\sim}\mathfrak{v}\\ \mathfrak{u}\neq\mathfrak{v}}}\frac{p_i(\mathfrak{v},\mathfrak{u})\sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t})}{p_i(\mathfrak{v},\mathfrak{v})}.$$

(3) If If $\mathfrak{t}^{(i-1)}\neq\mathfrak{t}^{(i)}$ *and* $\mathfrak{t}^{(i-\frac{1}{2})}\neq\mathfrak{t}^{(i+\frac{1}{2})}$*, and* $\mathfrak{t}\sigma_{i+\frac{1}{2}}$ *does not exist, then*

$$\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})=\frac{\delta_{\mathfrak{s}\mathfrak{t}}}{c_{\mathfrak{t}}(i+\frac{1}{2})-c_{\mathfrak{t}}(i-\frac{1}{2})}.$$

(4) If $\mathfrak{t}^{(i-1)}\neq\mathfrak{t}^{(i)}$ *and* $\mathfrak{t}^{(i-\frac{1}{2})}\neq\mathfrak{t}^{(i+\frac{1}{2})}$*, and* $\mathfrak{t}\sigma_{i+\frac{1}{2}}$ *exists, then*

$$\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})=\begin{cases}\dfrac{1}{c_{\mathfrak{t}}(i+\frac{1}{2})-c_{\mathfrak{t}}(i-\frac{1}{2})}, & \textit{if } \mathfrak{s}=\mathfrak{t},\\ 1-\dfrac{1}{(c_{\mathfrak{t}}(i+\frac{1}{2})-c_{\mathfrak{t}}(i-\frac{1}{2}))^2}, & \textit{if } \mathfrak{s}=\mathfrak{t}\sigma_{i+\frac{1}{2}} \textit{ and } \mathfrak{s}\succ\mathfrak{t},\\ 1, & \textit{if } \mathfrak{s}=\mathfrak{t}\sigma_{i+\frac{1}{2}} \textit{ and } \mathfrak{t}\succ\mathfrak{s},\\ 0, & \textit{otherwise.}\end{cases}\tag{5.2}$$

Theorem 5.3. *Let* $(\lambda,\ell)\in\hat{A}_{k+1}$ *and* $\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+1}$*. If* $i=1,\dots,k$*, then the following statements hold.*

(1) If $\mathfrak{t}^{(i-\frac{1}{2})}=\mathfrak{t}^{(i+\frac{1}{2})}$*, and* $\mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$ *satisfies* $\mathfrak{v}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}$ *and* $\mathfrak{v}^{(i)}=\mathfrak{v}^{(i+1)}$*, then*

$$\sigma_{i+1}(\mathfrak{s},\mathfrak{t})=\frac{\delta_{\mathfrak{s}\mathfrak{t}}+(p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{v})c_{\mathfrak{v}}(i)p_{i+1}(\mathfrak{v},\mathfrak{v})-c_{\mathfrak{t}}(i))p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{v})-p_{i+1}(\mathfrak{s},\mathfrak{v})p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})}{c_{\mathfrak{s}}(i+1)-c_{\mathfrak{t}}(i)},\quad \textit{if } \mathfrak{s}\neq\mathfrak{v},$$

and

$$\sigma_{i+1}(\mathfrak{v},\mathfrak{t})=\frac{p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})}{p_{i+1}(\mathfrak{v},\mathfrak{v})p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{v})}c_{\mathfrak{t}}(i)-\sum_{\substack{\mathfrak{u}\overset{i+1}{\sim}\mathfrak{v}\\ \mathfrak{u}\neq\mathfrak{v}}}\frac{p_{i+1}(\mathfrak{v},\mathfrak{u})\sigma_{i+1}(\mathfrak{u},\mathfrak{t})}{p_{i+1}(\mathfrak{v},\mathfrak{v})}.$$

(2) If $\mathfrak{t}^{(i)}=\mathfrak{t}^{(i+1)}$ *and* $\mathfrak{t}^{(i-\frac{1}{2})}\neq\mathfrak{t}^{(i+\frac{1}{2})}$*, and* $\mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$ *satisfies* $\mathfrak{v}\overset{i+1}{\sim}\mathfrak{t}$ *and* $\mathfrak{v}^{(i-\frac{1}{2})}=\mathfrak{v}^{(i+\frac{1}{2})}$*, then*

$$\sigma_{i+1}(\mathfrak{s},\mathfrak{t})=\frac{\delta_{\mathfrak{s}\mathfrak{t}}}{c_{\mathfrak{t}}(i+1)-c_{\mathfrak{t}}(i)},\qquad \textit{if } \mathfrak{s}\neq\mathfrak{v},$$

and

$$\sigma_{i+1}(\mathfrak{v},\mathfrak{t}) = \frac{p_{i+1}(\mathfrak{v},\mathfrak{t})}{(c_{\mathfrak{t}}(i)-c_{\mathfrak{t}}(i+1))p_{i+1}(\mathfrak{v},\mathfrak{v})}$$

(3) If $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+\frac{1}{2})}$ *and* $\mathfrak{t}^{(i)} \neq \mathfrak{t}^{(i+1)}$*, and* $\mathfrak{t}\sigma_{i+1}$ *does not exist, then*

$$\sigma_{i+1}(\mathfrak{s},\mathfrak{t}) = \frac{\delta_{\mathfrak{s}\mathfrak{t}}}{c_{\mathfrak{t}}(i+1)-c_{\mathfrak{t}}(i)}.$$

(4) If $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+\frac{1}{2})}$ *and* $\mathfrak{t}^{(i)} \neq \mathfrak{t}^{(i+1)}$*, and* $\mathfrak{t}\sigma_{i+1}$ *exists, then*

$$\sigma_{i+1}(\mathfrak{s},\mathfrak{t}) = \begin{cases} \dfrac{1}{c_{\mathfrak{t}}(i+1)-c_{\mathfrak{t}}(i)}, & \text{if } \mathfrak{s}=\mathfrak{t}, \\ 1-\dfrac{1}{(c_{\mathfrak{t}}(i+1)-c_{\mathfrak{t}}(i))^2}, & \text{if } \mathfrak{s}=\mathfrak{t}\sigma_{i+1} \text{ and } \mathfrak{s}\succ\mathfrak{t}, \\ 1, & \text{if } \mathfrak{s}=\mathfrak{t}\sigma_{i+1} \text{ and } \mathfrak{t}\succ\mathfrak{s}, \\ 0, & \text{otherwise.} \end{cases} \tag{5.3}$$

Remark 5.4. A consistent choice of signs for the square roots in (5.1) can be made as follows. Let $\mathfrak{s} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$. Suppose that for each $\mathfrak{t},\mathfrak{r} \in A^{\lambda}_{k+\frac{1}{2}}$, such that $\mathfrak{s} \overset{i}{\sim} \mathfrak{t}$ and $\mathfrak{s} \overset{i+\frac{1}{2}}{\sim} \mathfrak{r}$, a choice of scalars $\hat{p}_i(\mathfrak{t},\mathfrak{s})$ and $\hat{p}_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{s})$ has been made, subject to the constraints

$$\hat{p}_i(\mathfrak{t},\mathfrak{t}) = p_i(\mathfrak{t},\mathfrak{t}) \qquad \text{and} \qquad \hat{p}_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{r}) = p_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{r}), \tag{5.4}$$

and

$$\hat{p}_i(\mathfrak{t},\mathfrak{s})^2 = \frac{p_i(\mathfrak{t},\mathfrak{t})p_i(\mathfrak{s},\mathfrak{s})\langle f_{\mathfrak{s}}, f_{\mathfrak{s}}\rangle}{\langle f_{\mathfrak{t}}, f_{\mathfrak{t}}\rangle} \qquad \text{and} \qquad \hat{p}_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{s})^2 = \frac{p_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{r})p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{s})\langle f_{\mathfrak{s}}, f_{\mathfrak{s}}\rangle}{\langle f_{\mathfrak{r}}, f_{\mathfrak{r}}\rangle}. \tag{5.5}$$

If $\mathfrak{u},\mathfrak{v} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, and $\mathfrak{u} \overset{i}{\sim} \mathfrak{s}$ and $\mathfrak{v} \overset{i+\frac{1}{2}}{\sim} \mathfrak{s}$, determine uniquely the scalars $\hat{p}_i(\mathfrak{u},\mathfrak{t})$ and $\hat{p}_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{r})$ respectively by

$$\hat{p}_i(\mathfrak{u},\mathfrak{t})\hat{p}_i(\mathfrak{t},\mathfrak{s}) = \hat{p}_i(\mathfrak{u},\mathfrak{s})p_i(\mathfrak{t},\mathfrak{t}) \qquad \text{and} \qquad \hat{p}_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{r})\hat{p}_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{s}) = \hat{p}_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{s})p_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{r}). \tag{5.6}$$

Let $\hat{p}_i \in \mathrm{End}_{\kappa}\left(A^{(\lambda,\ell)}_{k+\frac{1}{2}}\right)$ and $\hat{p}_{i+\frac{1}{2}} \in \mathrm{End}_{\kappa}\left(A^{(\lambda,\ell)}_{k+\frac{1}{2}}\right)$ denote the matrices with entries as determined using (5.4),(5.5) and (5.6) above. Propositions 5.7 and 5.8 imply that, with the above choices, the maps

$$p_i \mapsto \hat{p}_i \qquad \text{and} \qquad p_{i+\frac{1}{2}} \mapsto \hat{p}_{i+\frac{1}{2}}, \qquad \text{for } i=1,\dots,k,$$

respect the relations (2)(i), (2)(ii), (3)(i)–(iii) and (4)(i), (4)(ii) in the presentation of Theorem 2.1, and hence yield an algebra homomorphism $\langle p_i, p_{i+\frac{1}{2}} \mid i=1,\dots,k\rangle \to \mathrm{End}_{\kappa}\left(A^{(\lambda,\ell)}_{k+\frac{1}{2}}\right)$.

5.1. **Proof of Main Results.** Before proving Theorems 5.1, 5.2 and 5.3, we establish some basic properties of the generators $p_i, p_{i+\frac{1}{2}}, \sigma_{i+\frac{1}{2}}, \sigma_{i+1}$.

Let $\mathfrak{s},\mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$. Write $\mathfrak{s} \overset{i}{\sim} \mathfrak{t}$ if

$$\mathfrak{s}^{(\ell)} = \mathfrak{t}^{(\ell)}, \qquad \text{for } i=1,\dots,k, \qquad \text{and} \qquad \mathfrak{s}^{(\ell-\frac{1}{2})} \neq \mathfrak{t}^{(\ell-\frac{1}{2})} \Rightarrow \ell = i,$$

and write $\mathfrak{s} \overset{i+\frac{1}{2}}{\sim} \mathfrak{t}$ if

$$\mathfrak{s}^{(\ell-\frac{1}{2})} = \mathfrak{t}^{(\ell-\frac{1}{2})}, \qquad \text{for } i=1,\dots,k, \qquad \text{and} \qquad \mathfrak{s}^{(\ell)} \neq \mathfrak{t}^{(\ell)} \Rightarrow \ell = i.$$

The next statement is obtained using Theorem 4.1 together with the commutativity relations of Proposition 2.4.

Proposition 5.5. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$ *and* $\mathfrak{s},\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$. *For* $i=1,\dots,k$, *the following statements hold:*

(1) If $p_i(\mathfrak{s},\mathfrak{t})\neq 0$, *then* $\mathfrak{s}\overset{i}{\sim}\mathfrak{t}$.

(2) If $p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})\neq 0$, *then* $\mathfrak{s}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}$.

Using Theorem 4.28 of [HR], or Schur–Weyl duality (*cf.* Theorem 3.6 of [HR] and Lemma 3.5 of [Na]), we obtain the next statement.

Proposition 5.6. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$ *and* $\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$. *Then, for* $i=1,\dots,k$, *the following statements hold:*

(1) If $\mathfrak{t}^{(i-\frac{1}{2})}=\mathfrak{t}^{(i+\frac{1}{2})}$ *then* $p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})\neq 0$.

(2) If $\mathfrak{t}^{(i)}=\mathfrak{t}^{(i+1)}$ *then* $p_{i+1}(\mathfrak{t},\mathfrak{t})\neq 0$.

The next statement provides a counterpart to Corollary 3.7 of [Na].

Proposition 5.7. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$. *Then, for* $i=1,\dots,k$, *the following equalities hold:*

$$p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{u})=p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{u}),\qquad \textit{for } \mathfrak{s},\mathfrak{t},\mathfrak{u}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}},\tag{5.7}$$

and

$$p_i(\mathfrak{s},\mathfrak{t})p_i(\mathfrak{t},\mathfrak{u})=p_i(\mathfrak{t},\mathfrak{t})p_i(\mathfrak{s},\mathfrak{u}),\qquad \textit{for } \mathfrak{s},\mathfrak{t},\mathfrak{u}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}.\tag{5.8}$$

Proof. It suffices to verify (5.7) for $i=k$. Let $\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$. If $\mathfrak{t}^{(k+\frac{1}{2})}\neq\mathfrak{t}^{(k-\frac{1}{2})}$, then

$$p_{k+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})=p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{s})=0,\qquad \text{for all } \mathfrak{s}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}},$$

so the relation (5.7) clearly holds. Suppose therefore, that $\mathfrak{t}^{(k+\frac{1}{2})}=\mathfrak{t}^{(k-\frac{1}{2})}$ so that

$$F_{\mathfrak{t}}p_{k+\frac{1}{2}}F_{\mathfrak{t}}=p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})F_{\mathfrak{t}}\neq 0.\tag{5.9}$$

Given $g(L_k,L_{k+\frac{1}{2}})$, a polynomial in L_k and $L_{k+\frac{1}{2}}$ over κ, then the Jones basic construction shows that

$$p_{k+\frac{1}{2}}g(L_k,L_{k+\frac{1}{2}})p_{k+\frac{1}{2}}=\xi p_{k+\frac{1}{2}}\qquad\text{where}\qquad \xi \text{ is a central element in } A_{k-\frac{1}{2}}(z).$$

In particular, if $\mathfrak{v}=\mathfrak{t}|_{k-\frac{1}{2}}$, then there exists a polynomial $g_{\mathfrak{t}}(L_k,L_{k+\frac{1}{2}})$ in $L_k,L_{k+\frac{1}{2}}$ over κ, such that

$$p_{k+\frac{1}{2}}F_{\mathfrak{t}}p_{k+\frac{1}{2}}=p_{k+\frac{1}{2}}g_{\mathfrak{t}}(L_k,L_{k+\frac{1}{2}})p_{k+\frac{1}{2}}F_{\mathfrak{v}}=p_{k+\frac{1}{2}}\xi_{\mathfrak{t}}F_{\mathfrak{v}}.\tag{5.10}$$

where $\xi_{\mathfrak{t}}$ is a central element in $A_{k-\frac{1}{2}}(z)$ and $F_{\mathfrak{v}}\in A_{k-\frac{1}{2}}$. Using (5.9) and (5.10),

$$F_{\mathfrak{t}}p_{k+\frac{1}{2}}F_{\mathfrak{t}}p_{k+\frac{1}{2}}F_{\mathfrak{t}}=\xi_{\mathfrak{t}}(F_{\mathfrak{t}}p_{k+\frac{1}{2}}F_{\mathfrak{t}})F_{\mathfrak{v}}=p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})\xi_{\mathfrak{t}}F_{\mathfrak{t}}F_{\mathfrak{v}}=p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})\xi_{\mathfrak{t}}F_{\mathfrak{t}},$$

and

$$F_{\mathfrak{t}}p_{k+\frac{1}{2}}F_{\mathfrak{t}}p_{k+\frac{1}{2}}F_{\mathfrak{t}}=p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})F_{\mathfrak{t}}p_{k+\frac{1}{2}}F_{\mathfrak{t}}=p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})^2F_{\mathfrak{t}}.$$

Since $F_{\mathfrak{t}}$ acts as a matrix unit on $A^{(\lambda,\ell)}_{k+\frac{1}{2}}$ and $p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})\neq 0$, it follows that

$$\xi_{\mathfrak{t}} \text{ acts on } f_{\mathfrak{v}}^{(k+\frac{1}{2})} \text{ by scalar multiplication by } p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t}).$$

Since $\xi_{\mathfrak{t}}$ is central in $A_{k-\frac{1}{2}}(z)$, it follows that

$$\xi_{\mathfrak{t}} \text{ acts on the } A_{k-\frac{1}{2}}\text{–module } A^{(\lambda,\ell-1)}_{k-\frac{1}{2}} \text{ by scalar multiplication by } p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{t}).$$

If $\mathfrak{s},\mathfrak{u}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, let $e_{\mathfrak{su}}$ denote the vector space endomorphism of $A^{(\lambda,\ell)}_{k+\frac{1}{2}}$ defined by

$$f^{(k+\frac{1}{2})}_{\mathfrak{r}}e_{\mathfrak{su}}=\delta_{\mathfrak{rs}}f^{(k+\frac{1}{2})}_{\mathfrak{u}}, \qquad\qquad \text{for } \mathfrak{r}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}.$$

Then, as operators on $A^{(\lambda,\ell)}_{k+\frac{1}{2}}$,

$$p_{k+\frac{1}{2}}F_{\mathfrak{t}}p_{k+\frac{1}{2}}=F_{\mathfrak{v}}\xi_{\mathfrak{t}}p_{k+\frac{1}{2}}\mathfrak{t}=\xi_{\mathfrak{t}}F_{\mathfrak{v}}p_{i+\frac{1}{2}}=p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})\sum_{\mathfrak{s}\overset{i+\frac{1}{2}}{\sim}\mathfrak{u}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}}p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{u})e_{\mathfrak{su}},$$

and

$$\begin{aligned}p_{k+\frac{1}{2}}F_{\mathfrak{t}}p_{k+\frac{1}{2}}=p_{k+\frac{1}{2}}F_{\mathfrak{t}}F_{\mathfrak{t}}p_{k+\frac{1}{2}}&=\Bigg(\sum_{\mathfrak{s}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}}p_{k+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})e_{\mathfrak{st}}\Bigg)\Bigg(\sum_{\mathfrak{u}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}}p_{k+\frac{1}{2}}(\mathfrak{t},\mathfrak{u})e_{\mathfrak{tu}}\Bigg)\\&=\sum_{\mathfrak{s}\overset{i+\frac{1}{2}}{\sim}\mathfrak{u}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}}p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{u})e_{\mathfrak{su}},\end{aligned}$$

which completes the proof of (5.9). The statement (5.10) follows by an appropriate modification of the proof of (5.9). □

Proposition 5.8 is a counterpart to Lemma 3.6 of [Na].

Proposition 5.8. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$ *and* $\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$.

(1) If $\mathfrak{t}^{(i-\frac{1}{2})}=\mathfrak{t}^{(i+\frac{1}{2})}$, *then*

$$f^{(k+\frac{1}{2})}_{\mathfrak{v}}p_{i+\frac{1}{2}}=\frac{p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{v})}{p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})}f^{(k+\frac{1}{2})}_{\mathfrak{t}}p_{i+\frac{1}{2}}, \qquad \textit{for all } \mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}} \textit{ such that } \mathfrak{v}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t},$$

and the eigenspace for the action of $p_{i+\frac{1}{2}}$ *on*

$$\mathrm{Span}_{\kappa}\{f^{(k+\frac{1}{2})}_{\mathfrak{s}}\mid\mathfrak{s}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}} \textit{ and } \mathfrak{s}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}\},$$

is one–dimensional.

(2) If $\mathfrak{t}^{(i-1)}=\mathfrak{t}^{(i)}$, *then*

$$f^{(k+\frac{1}{2})}_{\mathfrak{v}}p_{i}=\frac{p_{i}(\mathfrak{t},\mathfrak{v})}{p_{i}(\mathfrak{t},\mathfrak{t})}f^{(k+\frac{1}{2})}_{\mathfrak{t}}p_{i}, \qquad \textit{for all } \mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}} \textit{ such that } \mathfrak{v}\overset{i}{\sim}\mathfrak{t},$$

and the eigenspace for the action of p_i *on*

$$\mathrm{Span}_{\kappa}\{f^{(k+\frac{1}{2})}_{\mathfrak{s}}\mid\mathfrak{s}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}} \textit{ and } \mathfrak{s}\overset{i}{\sim}\mathfrak{t}\},$$

is one–dimensional.

Proof. (1) If $\mathfrak{s}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, then

$$p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{v})=p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{v})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})^{-1},$$

and

$$f^{(k+\frac{1}{2})}_{\mathfrak{v}}p_{i+\frac{1}{2}}=p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{v})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})^{-1}\sum_{\mathfrak{s}\overset{i+\frac{1}{2}}{\sim}\mathfrak{t}}p_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})f^{(k+\frac{1}{2})}_{\mathfrak{s}}=p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{v})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})^{-1}f_{\mathfrak{t}}p_{i+\frac{1}{2}},$$

as required. The proof of (2) is similar. □

Lemma 5.9. *Let* $(\lambda,\ell)\in\hat{A}_{k+\frac{1}{2}}$ *and* $\mathfrak{t}\in\hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$. *The following statements hold:*

(1) If $\mathfrak{t}^{(i-1)}\neq\mathfrak{t}^{(i)}$, *then* $c_{\mathfrak{t}}(i-\frac{1}{2})+c_{\mathfrak{t}}(i)\neq z$.

(2) If $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+\frac{1}{2})}$*, then* $c_{\mathfrak{t}}(i) + c_{\mathfrak{t}}(i+\frac{1}{2}) \neq z$*.*
(3) If $\mathfrak{t}^{(i-1)} \neq \mathfrak{t}^{(i)}$*, then* $f_{\mathfrak{t}} p_i = 0$*.*
(4) If $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+\frac{1}{2})}$*, then* $f_{\mathfrak{t}} p_{i+\frac{1}{2}} = 0$*.*

Proof. (1) Let $(\mathfrak{t}^{(i-1)}, \mathfrak{t}^{(i-\frac{1}{2})}, \mathfrak{t}^{(i-1)}) = (\mu, \nu, \upsilon)$. By definition, $c_{\mathfrak{t}}(i-\frac{1}{2}) + c_{\mathfrak{t}}(i) = z$ if and only if either $\nu = \upsilon$ and $\mu = \nu$, or there exists a node b such that $\upsilon = \nu \cup \{b\}$ and $\mu = \nu \cup \{b\}$. The proof of (2) is similar.
(3) Since the Jucys–Murphy elements act diagonally on the seminormal basis, we obtain

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}(L_{i-\frac{1}{2}} + L_i)p_i = (c_{\mathfrak{t}}(i-\tfrac{1}{2}) + c_{\mathfrak{t}}(i)) f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_i,$$

while, item (11) of Proposition 2.4 gives

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}(L_{i-\frac{1}{2}} + L_i)p_i = z f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_i.$$

The last statement is in contradiction to item (1) if $f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_i \neq 0$. The proof of (4) is similar. □

Let $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$ and $\mathfrak{s}, \mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$. Write $\mathfrak{s} \overset{i+1}{\simeq} \mathfrak{t}$ if

$$\mathfrak{s}^{(\ell+\frac{1}{2})} \neq \mathfrak{t}^{(\ell+\frac{1}{2})} \Rightarrow \ell = i \qquad \text{and} \qquad \mathfrak{s}^{(\ell)} \neq \mathfrak{t}^{(\ell)} \Rightarrow \ell = i,$$

and write $\mathfrak{s} \overset{i+\frac{1}{2}}{\simeq} \mathfrak{t}$ if

$$\mathfrak{s}^{(\ell-\frac{1}{2})} \neq \mathfrak{t}^{(\ell-\frac{1}{2})} \Rightarrow \ell = i \qquad \text{and} \qquad \mathfrak{s}^{(\ell)} \neq \mathfrak{t}^{(\ell)} \Rightarrow \ell = i.$$

The next statement is obtained using Theorem 4.1 together with the commutativity relations of Proposition 2.4.

Proposition 5.10. *Let* $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$ *and* $\mathfrak{s}, \mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$*. For* $i = 1, \dots, k$*, the following statements hold:*

(1) If $\sigma_{i+\frac{1}{2}}(\mathfrak{s}, \mathfrak{t}) \neq 0$*, then* $\mathfrak{s} \overset{i+\frac{1}{2}}{\simeq} \mathfrak{t}$*.*
(2) If $\sigma_{i+1}(\mathfrak{s}, \mathfrak{t}) \neq 0$*, then* $\mathfrak{s} \overset{i+1}{\simeq} \mathfrak{t}$*.*

If λ, μ are partitions, let $\lambda \ominus \mu = \lambda \setminus \mu \cup \mu \setminus \lambda$.

Lemma 5.11. *If* $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$ *and* $\mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$*, then the following statements hold:*

(1) If $\mathfrak{t}^{(i-1)} \neq \mathfrak{t}^{(i+\frac{1}{2})}$ *and the nodes* $\mathfrak{t}^{(i-1)} \ominus \mathfrak{t}^{(i+\frac{1}{2})}$ *are neither in the same row nor the same column, then there exists* $\mathfrak{s} = \mathfrak{t}\sigma_{i+\frac{1}{2}} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$*, uniquely determined by* $\mathfrak{s} \overset{i+\frac{1}{2}}{\simeq} \mathfrak{t}$ *and*

$$c_{\mathfrak{s}}(i+\tfrac{1}{2}) = c_{\mathfrak{t}}(i-\tfrac{1}{2}), \qquad c_{\mathfrak{s}}(i-\tfrac{1}{2}) = c_{\mathfrak{t}}(i+\tfrac{1}{2}), \qquad \textit{and}, \qquad c_{\mathfrak{s}}(i) = c_{\mathfrak{t}}(i).$$

(2) If $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+1)}$ *and the nodes* $\mathfrak{t}^{(i-\frac{1}{2})} \ominus \mathfrak{t}^{(i+1)}$ *are neither in the same row nor in the same column, then there exists* $\mathfrak{s} = \mathfrak{t}\sigma_{i+1} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$*, uniquely determined by* $\mathfrak{s} \overset{i+1}{\simeq} \mathfrak{t}$ *and*

$$c_{\mathfrak{s}}(i) = c_{\mathfrak{t}}(i+1), \qquad c_{\mathfrak{s}}(i+1) = c_{\mathfrak{t}}(i), \qquad \textit{and} \qquad c_{\mathfrak{s}}(i+\tfrac{1}{2}) = c_{\mathfrak{t}}(i+\tfrac{1}{2}).$$

Proof. (1) There is no loss of generality in proving the statement for $i = k$. We have two cases to consider.
(a) If $(\mathfrak{t}^{(k-1)}, \mathfrak{t}^{(k-\frac{1}{2})}, \mathfrak{t}^{(k)}, \mathfrak{t}^{(k+\frac{1}{2})}) = (\mu, \nu, \lambda, \lambda)$, where $\nu = \mu \setminus \{\alpha\} = \lambda \setminus \{\beta\}$ and α, β are neither

in the same row nor in the same column, then $\upsilon = \mu \cup \{\beta\} = \lambda \cup \{\alpha\}$ is a partition. Thus $\mathfrak{s} = \mathfrak{t}\sigma_{k+\frac{1}{2}} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$, given by

$$\mathfrak{s} \overset{k+\frac{1}{2}}{\simeq} \mathfrak{t} \qquad \text{and} \qquad (\mathfrak{s}^{(k-1)}, \mathfrak{s}^{(k-\frac{1}{2})}, \mathfrak{s}^{(k)}, \mathfrak{s}^{(k+\frac{1}{2})}) = (\mu, \mu, \upsilon, \lambda),$$

satisfies the required properties. By symmetry, we may write $\mathfrak{t} = \mathfrak{s}\sigma_{k+\frac{1}{2}}$.

(b) If $(\mathfrak{t}^{(k-1)}, \mathfrak{t}^{(k-\frac{1}{2})}, \mathfrak{t}^{(k)}, \mathfrak{t}^{(k+\frac{1}{2})}) = (\mu, \nu, \nu, \lambda)$, where $\nu = \mu \setminus \{\alpha\} = \lambda \cup \{\beta\}$ and α, β are neither in the same row nor in the same column, then $\upsilon = \mu \setminus \{\beta\} = \lambda \cup \{\alpha\}$ is a partition. Now define $\mathfrak{s} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$ by

$$\mathfrak{s} \overset{k+\frac{1}{2}}{\simeq} \mathfrak{t} \qquad \text{and} \qquad (\mathfrak{s}^{(k-1)}, \mathfrak{s}^{(k-\frac{1}{2})}, \mathfrak{s}^{(k)}, \mathfrak{s}^{(k+\frac{1}{2})}) = (\mu, \upsilon, \upsilon, \lambda)$$

to obtain the path in $\hat{A}$ with the required properties.

(2) There is no loss of generality supposing that $\mathfrak{t} \in A^{\lambda}_{k+1}$ and proving the statement for $i = k$. Again we have two cases to consider.

(a) If $(\mathfrak{t}^{(k-\frac{1}{2})}, \mathfrak{t}^{(k)}, \mathfrak{t}^{(k+\frac{1}{2})}, \mathfrak{t}^{(k+1)}) = (\mu, \nu, \lambda, \lambda)$, where $\nu = \mu \cup \{\alpha\} = \lambda \cup \{\beta\}$ and α, β are neither in the same row nor in the same column, then $\upsilon = \mu \setminus \{\beta\} = \lambda \setminus \{\alpha\}$ is a partition. Thus $\mathfrak{s} = \mathfrak{t}\sigma_{k+1} \in \hat{A}^{(\lambda,\ell)}_{k+1}$, given by

$$\mathfrak{s} \overset{k+1}{\simeq} \mathfrak{t} \qquad \text{and} \qquad (\mathfrak{s}^{(k-\frac{1}{2})}, \mathfrak{s}^{(k)}, \mathfrak{s}^{(k+\frac{1}{2})}, \mathfrak{s}^{(k+1)}) = (\mu, \mu, \upsilon, \lambda),$$

satisfies the required properties. By symmetry, we may define $\mathfrak{t} = \mathfrak{s}\sigma_{k+1}$.

(b) If $(\mathfrak{t}^{(k-\frac{1}{2})}, \mathfrak{t}^{(k)}, \mathfrak{t}^{(k+\frac{1}{2})}, \mathfrak{t}^{(k+1)}) = (\mu, \nu, \nu, \lambda)$, where $\nu = \mu \cup \{\alpha\} = \lambda \setminus \{\beta\}$ and α, β are neither in the same row nor in the same column, then $\upsilon = \mu \cup \{\beta\} = \lambda \setminus \{\alpha\}$ is a partition. Thus $\mathfrak{s} \in \hat{A}^{(\lambda,\ell)}_{k+1}$ defined by

$$\mathfrak{s} \overset{k+1}{\simeq} \mathfrak{t} \qquad \text{and} \qquad (\mathfrak{s}^{(k-\frac{1}{2})}, \mathfrak{s}^{(k)}, \mathfrak{s}^{(k+\frac{1}{2})}, \mathfrak{s}^{(k+1)}) = (\mu, \upsilon, \upsilon, \lambda).$$

satisfies the required properties. □

Proposition 5.12. *Let* $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$. *If* $\mathfrak{t} \in \hat{A}^{(\lambda,\ell)}_{k+\frac{1}{2}}$ *satisfies* $\mathfrak{t}^{(k-1)} = \mathfrak{t}^{(k-\frac{1}{2})} \subsetneq \mathfrak{t}^{(k)} = \mathfrak{t}^{(k+\frac{1}{2})}$, *then*

$$f^{(k+\frac{1}{2})}_{\mathfrak{t}^{\lambda}} p_{\mathfrak{t}} \sigma_{k+\frac{1}{2}} = f^{(k+\frac{1}{2})}_{\mathfrak{t}^{\lambda}} p_{\mathfrak{t}}.$$

Proof. Let $\mathfrak{t}^{(k-\frac{1}{2})} = \mu$, where $\lambda = \mu \cup \{(j, \lambda_\ell)\}$. Then,

$$p_{\mathfrak{t}^{(k)} \to \mathfrak{t}^{(k+\frac{1}{2})}} = p_{\mathfrak{t}^{(k-1)} \to \mathfrak{t}^{(k-\frac{1}{2})}} = 1, \quad \text{and} \quad p_{\mathfrak{t}^{(k-\frac{1}{2})} \to \mathfrak{t}^{(k)}} = w_{a_j,k}, \quad \text{where} \quad a_j = \ell + \sum_{r=1}^{j} \lambda_r,$$

If $j = 1, 2, \dots,$ define

$$\sigma^{(1)}_{j-\frac{1}{2}} = \sigma_{j-\frac{1}{2}} \qquad \text{and} \qquad \sigma^{(r+1)}_{j-\frac{1}{2}} = w_{r,r+j}\sigma^{(r)}_{j-\frac{1}{2}} w_{r+j,r} \qquad \text{for } r = 1, 2, \dots.$$

Since $\sigma_1 = \sigma_{1+\frac{1}{2}} = 1$, by induction we obtain

$$p_r \sigma^{(r)}_{j+\frac{1}{2}} = \begin{cases} p_r, & \text{if } j = 0 \text{ or } j = 1, \\ p_r \sigma^{(r+1)}_{j-\frac{1}{2}}, & \text{if } j \geqslant 2. \end{cases}$$

Thus, as $\sigma_{k-\ell+\frac{1}{2}} \mapsto 1$ under the map $\mathcal{A}_{k-\ell+\frac{1}{2}} \to \kappa\mathfrak{S}_{k-\ell}$ and $w_{a_j,k}$ commutes with $p^{(\ell)}_{\ell}$, we have

$$p^{(\ell)}_{\ell} w_{a_j,k} \sigma^{(\ell+1)}_{k-\ell+\frac{1}{2}} \equiv p^{(\ell)}_{\ell} w_{a_j,k} \mod A^{\rhd(\lambda,\ell)}_{k+\frac{1}{2}},$$

and

$$a_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_{\mathfrak{t}} \sigma_{k+\frac{1}{2}} \equiv a_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_{\mathfrak{t}} \mod A_{k+\frac{1}{2}}^{\rhd(\lambda,\ell)}.$$

Since $f_\lambda^{(k)} = \big(a_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})}\big)^* x_{(\lambda,\ell)}^{(k+\frac{1}{2})} a_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})}$, the statement follows. □

Proof of Theorem 5.1. The given formulae for the diagonal entries of the matrix representations p_i and $p_{i+\frac{1}{2}}$ follow from Proposition 6.2 and the expressions (6.19) and (6.18) respectively.

Let $\mathfrak{s}, \mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$. We prove our formula for the off–diagonal entry $p_i(\mathfrak{s},\mathfrak{t})$. If $p_i(\mathfrak{t},\mathfrak{t}) = 0$, then $p_i(\mathfrak{u},\mathfrak{t}) = 0$ for all $\mathfrak{u} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$, and the first equality in (5.1) clearly holds. Otherwise, by the contravariance of the bilinear form (4.1), we have

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_i F_{\mathfrak{s}\mathfrak{v}} = p_i(\mathfrak{s},\mathfrak{t}) \langle f_{\mathfrak{s}}^{(k+\frac{1}{2})}, f_{\mathfrak{s}}^{(k+\frac{1}{2})} \rangle f_{\mathfrak{v}}^{(k+\frac{1}{2})} \qquad \text{for all } \mathfrak{v} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)},$$

and

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_i F_{\mathfrak{s}\mathfrak{v}} = p_i(\mathfrak{t},\mathfrak{s}) \langle f_{\mathfrak{t}}^{(k+\frac{1}{2})}, f_{\mathfrak{t}}^{(k+\frac{1}{2})} \rangle f_{\mathfrak{v}} \qquad \text{for all } \mathfrak{v} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}.$$

By Proposition 5.7, we have $p_i(\mathfrak{t},\mathfrak{s}) = p_i(\mathfrak{s},\mathfrak{s}) p_i(\mathfrak{t},\mathfrak{t}) p_i(\mathfrak{s},\mathfrak{t})^{-1}$, and

$$p_i(\mathfrak{s},\mathfrak{t})^2 = \frac{p_i(\mathfrak{s},\mathfrak{s}) p_i(\mathfrak{t},\mathfrak{t}) \langle f_{\mathfrak{t}}^{(k+\frac{1}{2})}, f_{\mathfrak{t}}^{(k+\frac{1}{2})} \rangle}{\langle f_{\mathfrak{s}}^{(k+\frac{1}{2})}, f_{\mathfrak{s}}^{(k+\frac{1}{2})} \rangle},$$

as required. The proof of the second equality in (5.1) is similar. □

Proof of Theorem 5.2. Let $(\lambda,\ell) \in \hat{A}_{k+\frac{1}{2}}$ and $\mathfrak{s}, \mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$.

(1) Suppose that $\mathfrak{t}^{(i-\frac{1}{2})} = \mathfrak{t}^{(i+\frac{1}{2})}$ and let $\mathfrak{v} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ such that $\mathfrak{v} \overset{i+\frac{1}{2}}{\sim} \mathfrak{t}$ and $\mathfrak{v}^{(i-1)} = \mathfrak{v}^{(i)}$. If $\mathfrak{s} \neq \mathfrak{v}$, then, using item (2) of Proposition 2.5 and Lemma 5.9,

$$\begin{aligned}\sum_{\mathfrak{u} \overset{i+\frac{1}{2}}{\simeq} \mathfrak{t}} (c_{\mathfrak{u}}(i+\tfrac{1}{2}) - c_{\mathfrak{t}}(i-\tfrac{1}{2}) \sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t}) f_{\mathfrak{u}}^{(k+\frac{1}{2})} &= -f_{\mathfrak{t}}^{(k+\frac{1}{2})} (p_i p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} L_i - (z - L_{i-\frac{1}{2}}) p_{i+\frac{1}{2}} - 1) \\ &= f_{\mathfrak{t}}^{(k+\frac{1}{2})} - f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}} L_i - p_i(\mathfrak{v},\mathfrak{t}) f_{\mathfrak{v}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}} + (z - c_{\mathfrak{t}}(i-\tfrac{1}{2})) f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}},\end{aligned}$$

which gives the formula for $\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})$. Next,

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})} \sigma_{i+\frac{1}{2}} p_i p_{i+\frac{1}{2}} = \sum_{\mathfrak{u} \overset{i}{\sim} \mathfrak{v}} \sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t}) f_{\mathfrak{u}}^{(k+\frac{1}{2})} p_i p_{i+\frac{1}{2}} = \sum_{\mathfrak{u} \overset{i}{\sim} \mathfrak{v}} \sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t}) p_i(\mathfrak{v},\mathfrak{u}) f_{\mathfrak{v}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}},$$

where

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})} \sigma_{i+\frac{1}{2}} p_i p_{i+\frac{1}{2}} = c_{\mathfrak{t}}(i) f_{\mathfrak{t}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}} = c_{\mathfrak{t}}(i) p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t}) p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{v})^{-1} f_{\mathfrak{v}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}}$$

and $f_{\mathfrak{v}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}} \neq 0$. The formula for $\sigma_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})$ now follows.

(2) Suppose that $\mathfrak{t}^{(i-1)} = \mathfrak{t}^{(i)}$ and $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+\frac{1}{2})}$, and let $\mathfrak{v} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ such that $\mathfrak{v} \overset{i}{\sim} \mathfrak{t}$ and $\mathfrak{v}^{(i-\frac{1}{2})} = \mathfrak{v}^{(i+\frac{1}{2})}$. Using item (2) of Proposition 2.5 and Lemma 5.9,

$$\sum_{\mathfrak{u} \overset{i+\frac{1}{2}}{\simeq} \mathfrak{t}} (c_{\mathfrak{u}}(i+\tfrac{1}{2}) - c_{\mathfrak{t}}(i-\tfrac{1}{2}) \sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t}) f_{\mathfrak{u}}^{(k+\frac{1}{2})} = f_{\mathfrak{t}} - p_i(\mathfrak{v},\mathfrak{t}) f_{\mathfrak{v}}^{(k+\frac{1}{2})} p_{i+\frac{1}{2}}$$

which gives the formula for $\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})$ when $\mathfrak{s}\neq\mathfrak{v}$. Next,

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}\sigma_{i+\frac{1}{2}}p_ip_{i+\frac{1}{2}}=\sum_{\mathfrak{u}\overset{i}{\sim}\mathfrak{v}}\sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t})f_{\mathfrak{u}}^{(k+\frac{1}{2})}p_ip_{i+\frac{1}{2}}=\sum_{\mathfrak{u}\overset{i}{\sim}\mathfrak{v}}\sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t})p_i(\mathfrak{v},\mathfrak{u})f_{\mathfrak{v}}^{(k+\frac{1}{2})}p_{i+\frac{1}{2}},$$

where

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}\sigma_{i+\frac{1}{2}}p_ip_{i+\frac{1}{2}}=c_{\mathfrak{t}}(i)f_{\mathfrak{t}}^{(k+\frac{1}{2})}p_{i+\frac{1}{2}}=0\qquad\text{and}\qquad f_{\mathfrak{v}}^{(k+\frac{1}{2})}p_{i+\frac{1}{2}}\neq 0.$$

The formula for $\sigma_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})$ now follows.

(3) Suppose that $\mathfrak{t}^{(i-1)}\neq\mathfrak{t}^{(i)}$ and $\mathfrak{t}^{(i-\frac{1}{2})}\neq\mathfrak{t}^{(i+\frac{1}{2})}$ and that $\mathfrak{t}\sigma_{i+\frac{1}{2}}$ does not exist. By item (2) of Proposition 2.5 and Lemma 5.9,

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}\sigma_{i+\frac{1}{2}}L_{i+\frac{1}{2}}-f_{\mathfrak{t}}^{(k+\frac{1}{2})}L_{i-\frac{1}{2}}\sigma_{i+\frac{1}{2}}=f_{\mathfrak{t}}^{(k+\frac{1}{2})}.$$

Since

$$\mathfrak{u}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}\text{ and }\mathfrak{u}\overset{i+\frac{1}{2}}{\simeq}\mathfrak{t}\qquad\text{implies that}\qquad\mathfrak{u}=\mathfrak{t},$$

it follows that $\sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{t})=0$ if $\mathfrak{u}\neq\mathfrak{t}$, while

$$\big(c_{\mathfrak{t}}(i+\tfrac{1}{2})-c_{\mathfrak{t}}(i-\tfrac{1}{2})\big)\,\sigma_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})=1,$$

as required.

(4) Suppose that $\mathfrak{t}^{(i-1)}\neq\mathfrak{t}^{(i)}$ and $\mathfrak{t}^{(i-\frac{1}{2})}\neq\mathfrak{t}^{(i+\frac{1}{2})}$, and that $\mathfrak{t}\sigma_{i+\frac{1}{2}}$ exists. First observe that the hypotheses on $\mathfrak{t}$ imply that if $\mathfrak{u}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ and $\mathfrak{u}\overset{i+\frac{1}{2}}{\simeq}\mathfrak{t}$, then $\mathfrak{u}\in\{\mathfrak{t},\mathfrak{t}\sigma_{i+\frac{1}{2}}\}$. Thus, by Lemma 5.9,

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}\sigma_{i+\frac{1}{2}}=\sigma_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})f_{\mathfrak{t}}^{(k+\frac{1}{2})}+\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})f_{\mathfrak{s}}^{(k+\frac{1}{2})},\qquad\text{where }\mathfrak{s}=\mathfrak{t}\sigma_{i+\frac{1}{2}}.$$

We now determine $\sigma_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})$. Lemma 5.9 and the assumptions on $\mathfrak{t}$ imply that

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}p_i=f_{\mathfrak{t}}^{(k+\frac{1}{2})}p_{i+\frac{1}{2}}=0.$$

Thus item (2) of Proposition 2.5 yields

$$(c_{\mathfrak{t}}(i+\tfrac{1}{2})-c_{\mathfrak{t}}(i-\tfrac{1}{2}))\sigma_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})=1,$$

which gives the required expression for $\sigma_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})$.

Next, we show that if $\mathfrak{s}=\mathfrak{t}\sigma_{i+\frac{1}{2}}$ and $\mathfrak{t}\succ\mathfrak{s}$, then $\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})=1$. Since

$$f_{\mathfrak{t}}^{(k+\frac{1}{2})}\sigma_{i+\frac{1}{2}}=f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})}p_{\mathfrak{t}}\sigma_{i+\frac{1}{2}}+\sum_{\mathfrak{u}\succ\mathfrak{t}}r_{\mathfrak{u}}f_{\mathfrak{u}}^{(k+\frac{1}{2})}\sigma_{i+\frac{1}{2}},\qquad\text{where }r_{\mathfrak{u}}\in\kappa,\text{ for }\mathfrak{u}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)},$$

it suffices to show that

$$\sigma_{i+\frac{1}{2}}(\mathfrak{u},\mathfrak{s})=0,\qquad\text{if }\mathfrak{u}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}\text{ and }\mathfrak{u}\succ\mathfrak{t},\tag{5.11}$$

and that there exist $r_{\mathfrak{u}}\in\kappa$, for $\mathfrak{u}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$, such that

$$f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})}p_{\mathfrak{t}}\sigma_{i+\frac{1}{2}}=f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})}p_{\mathfrak{s}}+\sum_{\mathfrak{u}\succ\mathfrak{s}}r_{\mathfrak{u}}f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})}p_{\mathfrak{u}}.\tag{5.12}$$

To prove (5.11), suppose that $\mathfrak{u}\succ\mathfrak{t}$. Then

$$f_{\mathfrak{u}}^{(k+\frac{1}{2})}\sigma_{i+\frac{1}{2}}=\sum_{\mathfrak{r}\overset{i+\frac{1}{2}}{\simeq}\mathfrak{u}}\sigma_{i+\frac{1}{2}}(\mathfrak{r},\mathfrak{u})f_{\mathfrak{r}}^{(k+\frac{1}{2})},\qquad\text{for }\mathfrak{r}\in\hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}.\tag{5.13}$$

If $\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{u}) \neq 0$ in the expression (5.13), then $\mathfrak{u} \overset{i+\frac{1}{2}}{\simeq} \mathfrak{s}$, so $\mathfrak{u} \in \{\mathfrak{t},\mathfrak{s}\}$, which leads to a contradiction if $\mathfrak{u} \succ \mathfrak{t} \succ \mathfrak{s}$. Thus (5.11) holds.

Now we verify (5.12) assuming, without any loss of generality, that $i = k$. There are two cases to consider here.

(a) Suppose that $\mathfrak{t}^{(i-\frac{1}{2})} \subsetneq \mathfrak{t}^{(i+\frac{1}{2})}$. Let

$$\big(\mathfrak{t}^{(k-1)}, \mathfrak{t}^{(k-\frac{1}{2})}, \mathfrak{t}^{(k)}, \mathfrak{t}^{(k+\frac{1}{2})}\big) = (\nu,\mu,\lambda,\lambda) \quad \text{and} \quad \big(\mathfrak{s}^{(k-1)}, \mathfrak{s}^{(k-\frac{1}{2})}, \mathfrak{s}^{(k)}, \mathfrak{s}^{(k+\frac{1}{2})}\big) = (\nu,\nu,\upsilon,\lambda),$$

where

$$\lambda = \mu \cup \{(j,\lambda_j)\} \quad \text{and} \quad \nu = \mu \cup \{(j',\nu_{j'})\} \quad \text{and} \quad \upsilon = \nu \cup \{(j,\lambda_j)\} = \lambda \cup \{(j',\nu_{j'})\}.$$

Then

$$p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}} = 1 \qquad \text{and} \qquad p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}} = w_{a_j,k}, \qquad \text{where} \qquad a_j = \ell + \sum_{r=1}^{j} \lambda_j,$$

and

$$p_{\mathfrak{t}^{(k-1)}\to\mathfrak{t}^{(k-\frac{1}{2})}} = w_{\ell,k-1} p_{k-\frac{1}{2}} w_{k-1,a_{j'}} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r}, \qquad \text{where} \qquad a_{j'} = \ell - 1 + \sum_{r=1}^{j'} \nu_r,$$

while

$$p_{\mathfrak{s}^{(k)}\to\mathfrak{s}^{(k+\frac{1}{2})}} = w_{\ell,k} p_{k+\frac{1}{2}} w_{k,\alpha_{j'}} \sum_{r=0}^{\lambda_{j'}} w_{\alpha_{j'},\alpha_{j'}-r}, \qquad \text{where} \qquad \alpha_{j'} = \ell - 1 + \sum_{r=1}^{j'} \upsilon_r,$$

and

$$p_{\mathfrak{s}^{(k-\frac{1}{2})}\to\mathfrak{s}^{(k)}} = w_{\alpha_j,k}, \qquad \text{and} \qquad p_{\mathfrak{s}^{(k-1)}\to\mathfrak{s}^{(k-\frac{1}{2})}} = 1, \qquad \text{where} \qquad \alpha_j = \ell - 1 + \sum_{r=1}^{j} \upsilon_r.$$

Therefore,

$$\begin{aligned} f_\lambda^{(k)} & p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}} p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}} p_{\mathfrak{t}^{(k-1)}\to\mathfrak{t}^{(k-\frac{1}{2})}} \sigma_{k+\frac{1}{2}} \\ &= f_\lambda^{(k)} w_{a_j,k} w_{\ell,k-1} p_{k-\frac{1}{2}} w_{k-1,a_{j'}} \Bigg(\sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r} \Bigg) \sigma_{k+\frac{1}{2}} \\ &= f_\lambda^{(k)} w_{a_j,k} w_{\ell,k-1} \sigma_k p_{k+\frac{1}{2}} w_{k,a_{j'}} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r}. \end{aligned}$$

Now, $f_\lambda^{(k)} w_{a_j,k} w_{\ell,k-1} = f_\lambda^{(k)} p_{\mathfrak{u}}$, where $\mathfrak{u} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ is given by

$$\big(\mathfrak{u}^{(k-\frac{3}{2})}, \mathfrak{u}^{(k-1)}, \mathfrak{u}^{(k-\frac{1}{2})}, \mathfrak{u}^{(k)}, \mathfrak{u}^{(k+\frac{1}{2})}\big) = (\mu,\mu,\mu,\lambda,\lambda),$$

and

$$p_{\mathfrak{u}^{(\frac{1}{2})}\to\mathfrak{u}^{(1)}} = p_{\mathfrak{u}^{(1)}\to\mathfrak{u}^{(1+\frac{1}{2})}} = \cdots = p_{\mathfrak{u}^{(k-2)}\to\mathfrak{u}^{(k-\frac{3}{2})}} = 1.$$

Thus, by Proposition 5.12,

$$f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_{\mathfrak{u}} \sigma_{k+\frac{1}{2}} = f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_{\mathfrak{u}},$$

and

$$f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{a_j,k} w_{\ell,k-1} \sigma_k p_{k+\frac{1}{2}} w_{k,a_{j'}} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r} = f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{a_j,k} w_{\ell,k-1} s_{k-1} p_{k+\frac{1}{2}} w_{k,a_{j'}} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r}.$$

There are two further cases to consider here.
(i) If $j < j'$, then, by the braid relation, the right hand side of the last expression is

$$\begin{aligned} f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{a_j,k} w_{\ell,k} p_{k+\frac{1}{2}} w_{k,a_{j'}} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r} &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{\ell,k} p_{k+\frac{1}{2}} w_{a_j-1,k-1} w_{k,a_{j'}} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r} \\ &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{\ell,k} p_{k+\frac{1}{2}} w_{k,a_{j'}+1} w_{a_j-1,k} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r} \\ &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{\ell,k} p_{k+\frac{1}{2}} w_{k,a_{j'}+1} \left(\sum_{r=0}^{\mu_{j'}} w_{a_{j'}+1,a_{j'}-r+1} \right) w_{a_j-1,k} \\ &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{\ell,k} p_{k+\frac{1}{2}} w_{k,\alpha_{j'}} \left(\sum_{r=0}^{\lambda_{j'}} w_{\alpha_{j'},\alpha_{j'}-r} \right) w_{\alpha_j,k} \\ &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_{\mathfrak{s}^{(k)} \to \mathfrak{s}^{(k+\frac{1}{2})}} p_{\mathfrak{s}^{(k-\frac{1}{2})} \to \mathfrak{s}^{(k)}} p_{\mathfrak{s}^{(k-1)} \to \mathfrak{s}^{(k-\frac{1}{2})}}, \end{aligned}$$

since, in this case, $a_{j'} = \alpha_{j'} - 1$ and $w_{a_j-1,k-1} w_{k,a_{j'}} = w_{k,a_{j'}+1} w_{a_j-1,k}$.
(ii) If $j' < j$, then using the fact that in this case $a_j = \alpha_j$ and $a_{j'} = \alpha_{j'}$,

$$\begin{aligned} f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{a_j,k} w_{\ell,k} p_{k+\frac{1}{2}} w_{k,a_{j'}} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r} &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{\ell,k} p_{k+\frac{1}{2}} w_{k,a_{j'}} w_{a_j,k} \sum_{r=0}^{\mu_{j'}} w_{a_{j'},a_{j'}-r} \\ &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} w_{\ell,k} p_{k+\frac{1}{2}} w_{k,a_{j'}} \left(\sum_{r=0}^{\lambda_j} w_{a_{j'},a_{j'}-r} \right) w_{a_j,k} \\ &= f_{\mathfrak{t}^\lambda}^{(k+\frac{1}{2})} p_{\mathfrak{s}^{(k)} \to \mathfrak{s}^{(k+\frac{1}{2})}} p_{\mathfrak{s}^{(k-\frac{1}{2})} \to \mathfrak{s}^{(k)}} p_{\mathfrak{s}^{(k-1)} \to \mathfrak{s}^{(k-\frac{1}{2})}}. \end{aligned}$$

(b) Suppose that $\mathfrak{t}^{(k-\frac{1}{2})} \supsetneq \mathfrak{t}^{(k+\frac{1}{2})}$. We embed $A_{k+\frac{1}{2}}^{(\lambda,\ell)}$ in $A_{k+\frac{1}{2}}^{(\lambda,\ell+1)}$ and let

$$(\mathfrak{t}^{(k-1)}, \mathfrak{t}^{(k-\frac{1}{2})}, \mathfrak{t}^{(k)}, \mathfrak{t}^{(k+\frac{1}{2})}, \mathfrak{t}^{(k+1)}) = (\mu, \nu, \nu, \lambda, \lambda),$$

where $\mu \supsetneq \lambda$ and $\mu \setminus \lambda$ are neither in the same row, nor in the same column. The observe that, as in (3) of Theorem 5.3, $f_{\mathfrak{t}}^{(k+1)} \sigma_{k+1} = f_{\mathfrak{t}}^{(k+1)}$. Since $\sigma_{k+1} \sigma_{k+\frac{1}{2}} = s_k$, we have

$$\sigma_{i+\frac{1}{2}}(\mathfrak{s}, \mathfrak{t}) = s_k(\mathfrak{s}, \mathfrak{t}) \qquad \text{for all } \mathfrak{s} \in \hat{A}_{k+1}^{(\lambda,\ell)}.$$

Let us suppose that $\mathfrak{s} = \mathfrak{t}\sigma_{k+\frac{1}{2}}$, where $\mathfrak{t} \succ \mathfrak{s}$. Since

$$f_{\mathfrak{t}}^{(k+1)} s_k = f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{t}} s_k + \sum_{\mathfrak{u} \in \hat{A}_{k+1}^{(\lambda,\ell)}} r_{\mathfrak{u}} f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{u}} s_k, \qquad \text{where } a_{\mathfrak{u}} = 0 \text{ unless } \mathfrak{u} \succ \mathfrak{t},$$

it suffices to show that

$$s_k(\mathfrak{u}, \mathfrak{s}) = 0 \quad \text{if } \mathfrak{u} \in \hat{A}_{k+1}^{(\lambda,\ell)} \text{ and } \mathfrak{u} \succ \mathfrak{t}, \qquad \text{and} \qquad f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{t}} s_k = f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{s}}. \tag{5.14}$$

For the first equality in (5.14), observe that if $s_k(\mathfrak{u}, \mathfrak{s}) \neq 0$, then $\mathfrak{u}^{(k-1)} = \mathfrak{s}^{(k-1)}$ which, together with $\mathfrak{u}^{(k+1)} = \mathfrak{s}^{(k+1)}$, implies that $\mathfrak{u}^{(k+\frac{1}{2})} = \mathfrak{s}^{(k+\frac{1}{2})}$. Thus $\mathfrak{u} \in \{\mathfrak{s}, \mathfrak{t}\}$, which is impossible under the assumption that $\mathfrak{u} \succ \mathfrak{t} \succ \mathfrak{s}$.

For the second equality in (5.14), we define $\mathfrak{s} \in \hat{A}_{k+1}^{(\lambda,\ell)}$ by

$$p_{\mathfrak{s}^{(\frac{1}{2})} \to \mathfrak{s}^{(1)}} = p_{\mathfrak{t}^{(\frac{1}{2})} \to \mathfrak{t}^{(1)}}, p_{\mathfrak{s}^{(1)} \to \mathfrak{s}^{(1+\frac{1}{2})}} = p_{\mathfrak{t}^{(1)} \to \mathfrak{t}^{(1+\frac{1}{2})}}, \dots, p_{\mathfrak{s}^{(k+\frac{3}{2})} \to \mathfrak{s}^{(k-1)}} = p_{\mathfrak{t}^{(k+\frac{3}{2})} \to \mathfrak{t}^{(k-1)}},$$

and

$$(\mathfrak{s}^{(k-1)}, \mathfrak{s}^{(k-\frac{1}{2})}, \mathfrak{s}^{(k)}, \mathfrak{s}^{(k+\frac{1}{2})}, \mathfrak{s}^{(k+1)}) = (\mu, \upsilon, \upsilon, \lambda, \lambda),$$

where

$$\nu = \lambda \cup \{(j, \nu_j)\} \qquad \text{and} \qquad \mu = \nu \cup \{(j', \upsilon_{j'})\} = \upsilon \cup \{(j, \nu_j)\},$$

and $j < j'$, so that $\mathfrak{s} = \mathfrak{t}\sigma_{k+\frac{1}{2}}$ and $\mathfrak{t} \succ \mathfrak{s}$. If $\lambda \vdash k + 1 - \ell$, then

$$p_{\mathfrak{t}^{(k+\frac{1}{2})} \to \mathfrak{t}^{(k+1)}} = p_{\mathfrak{s}^{(k+\frac{1}{2})} \to \mathfrak{s}^{(k+1)}} = w_{\ell,k+1}, \qquad \text{and} \qquad p_{\mathfrak{t}^{(k-\frac{1}{2})} \to \mathfrak{t}^{(k)}} = p_{\mathfrak{s}^{(k-\frac{1}{2})} \to \mathfrak{s}^{(k)}} = w_{\ell-2,k},$$

and

$$p_{\mathfrak{t}^{(k)} \to \mathfrak{t}^{(k+\frac{1}{2})}} = w_{\ell-1,k} p_{k+\frac{1}{2}} w_{k,a_j} \sum_{r=0}^{\lambda_j} w_{a_j, a_j - r}, \qquad \text{where} \qquad a_j = \ell - 2 + \sum_{r=1}^{j} \nu_r,$$

and

$$p_{\mathfrak{t}^{(k-1)} \to \mathfrak{t}^{(k-\frac{1}{2})}} = w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-1,a_{j'}} \sum_{r=0}^{\nu_{j'}} w_{a_{j'}, a_{j'} - r}, \qquad \text{where} \qquad a_{j'} = \ell - 4 + \sum_{r=1}^{j'} \mu_r,$$

while

$$p_{\mathfrak{s}^{(k)} \to \mathfrak{s}^{(k+\frac{1}{2})}} = w_{\ell-1,k} p_{k+\frac{1}{2}} w_{k,\alpha_{j'}} \sum_{r=0}^{\lambda_{j'}} w_{\alpha_{j'}, \alpha_{j'} - r}, \qquad \text{where} \qquad \alpha_{j'} = \ell - 2 + \sum_{r=1}^{j'} \upsilon_r = a_{j'} + 1,$$

and

$$p_{\mathfrak{s}^{(k-1)} \to \mathfrak{s}^{(k-\frac{1}{2})}} = w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-1,\alpha_j} \sum_{r=0}^{\upsilon_j} w_{\alpha_j, \alpha_j - r}, \qquad \text{where} \qquad \alpha_j = \ell - 4 + \sum_{r=1}^{j} \mu_r = a_j - 2.$$

By the braid relation,

$$\begin{aligned} f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{t}^{(k+\frac{1}{2})} \to \mathfrak{t}^{(k+1)}} p_{\mathfrak{t}^{(k)} \to \mathfrak{t}^{(k+\frac{1}{2})}} p_{\mathfrak{t}^{(k-\frac{1}{2})} \to \mathfrak{t}^{(k)}} p_{\mathfrak{t}^{(k-1)} \to \mathfrak{t}^{(k-\frac{1}{2})}} s_k &= f_{\mathfrak{t}^\lambda}^{(k+1)} w_{\ell,k+1} w_{\ell-1,k} p_{k+\frac{1}{2}} w_{k,a_j} \\ &\quad \times w_{\ell-2,k} w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-1,a_{j'}} s_k \left(\sum_{r=0}^{\upsilon_j} w_{\alpha_j, \alpha_j - r} \right) \left(\sum_{r=0}^{\nu_{j'}} w_{a_{j'}, a_{j'} - r} \right), \end{aligned}$$

and

$$\begin{aligned} f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{s}^{(k)} \to \mathfrak{s}^{(k+\frac{1}{2})}} p_{\mathfrak{s}^{(k-\frac{1}{2})} \to \mathfrak{s}^{(k)}} p_{\mathfrak{s}^{(k-1)} \to \mathfrak{s}^{(k-\frac{1}{2})}} &= f_{\mathfrak{t}^\lambda}^{(k+1)} w_{\ell,k+1} w_{\ell-1,k} p_{k+\frac{1}{2}} w_{k,\alpha_{j'}} \\ &\quad \times w_{\ell-2,k} w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-1,\alpha_j} \left(\sum_{r=0}^{\nu_{j'}} w_{a_{j'}, a_{j'} - r} \right) \left(\sum_{r=0}^{\upsilon_j} w_{\alpha_j, \alpha_j - r} \right), \end{aligned}$$

so the fact that $f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{t}} s_k = f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{s}}$ will follow once we have shown that

$$p_{k+\frac{1}{2}} w_{k,a_j} w_{\ell-2,k} w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-1,a_{j'}} s_k = p_{k+\frac{1}{2}} w_{k,\alpha_{j'}} w_{\ell-2,k} w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-1,\alpha_j}. \tag{5.15}$$

Considering the left hand side of (5.15), the braid relation gives

$$\begin{aligned} p_{k+\frac{1}{2}} w_{k,a_j} w_{\ell-2,k} w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-1,a_{j'}} s_k &= p_{k+\frac{1}{2}} w_{\ell-2,k} w_{\ell-3,k-1} p_{k-\frac{1}{2}} w_{k-2,a_j-2} w_{k-1,a_{j'}} s_k \\ &= w_{\ell-2,k-1} w_{\ell-3,k-2} p_{k+\frac{1}{2}} s_{k-1} s_{k-2} p_{k-\frac{1}{2}} w_{k-2,a_j-2} w_{k-1,a_{j'}} s_k \\ &= w_{\ell-2,k-1} w_{\ell-3,k-2} p_{k-\frac{3}{2}} p_{k+\frac{1}{2}} s_{k-1} s_{k-2} s_k w_{k-2,a_j-2} w_{k-1,a_{j'}} \\ &= w_{\ell-2,k-1} w_{\ell-3,k-2} p_{k-\frac{3}{2}} p_{k+\frac{1}{2}} s_{k-1} w_{k-2,a_j-2} w_{k-1,a_{j'}}, \end{aligned}$$

since $p_{i-\frac{1}{2}}p_{i+\frac{3}{2}} = p_{i-\frac{1}{2}}p_{i+\frac{3}{2}}s_is_{i-1}s_{i+1}s_i$. Considering the right hand side of (5.15), the braid relation gives

$$\begin{aligned}p_{k+\frac{1}{2}}w_{k,\alpha_{j'}}w_{\ell-2,k}w_{\ell-3,k-1}p_{k-\frac{1}{2}}w_{k-1,\alpha_j} &= p_{k+\frac{1}{2}}w_{\ell-2,k}w_{\ell-3,k-1}p_{k-\frac{1}{2}}w_{k-2,\alpha_{j'}-2}w_{k-1,\alpha_j}\\ &= w_{\ell-2,k-1}w_{\ell-3,k-2}p_{k+\frac{1}{2}}s_{k-1}s_{k-2}p_{k-\frac{1}{2}}w_{k-1,\alpha_j}w_{k-1,\alpha_{j'}-1}\\ &= w_{\ell-2,k-1}w_{\ell-3,k-2}p_{k-\frac{3}{2}}p_{k+\frac{1}{2}}s_{k-1}w_{k-2,\alpha_j}w_{k-1,\alpha_{j'}-1},\end{aligned}$$

which completes the proof of (5.15). This verifies (5.12) and establishes that $\sigma_{k+\frac{1}{2}}(\mathfrak{s},\mathfrak{t}) = 1$ if $\mathfrak{s} = \mathfrak{t}\sigma_{k+\frac{1}{2}}$ and $\mathfrak{t} \succ \mathfrak{s}$.

Finally, using the above calculations together with the fact that $\sigma_{i+\frac{1}{2}}^2 = 1$, we can verify the given formula for $\sigma_{i+\frac{1}{2}}(\mathfrak{s},\mathfrak{t})$ when $\mathfrak{s} = \mathfrak{t}\sigma_{i+\frac{1}{2}}$ and $\mathfrak{s} \succ \mathfrak{t}$ by observing that $c_{\mathfrak{s}}(i-\frac{1}{2}) = c_{\mathfrak{t}}(i+\frac{1}{2})$ and $c_{\mathfrak{s}}(i+\frac{1}{2}) = c_{\mathfrak{t}}(i-\frac{1}{2})$. □

Proof of Theorem 5.3. Let $(\lambda,\ell) \in \hat{A}_{k+1}$ and $\mathfrak{s},\mathfrak{t} \in \hat{A}_{k+1}^{(\lambda,\ell)}$.

(1) Suppose that $\mathfrak{t}^{(i-\frac{1}{2})} = \mathfrak{t}^{(i+\frac{1}{2})}$ and let $\mathfrak{v} \in \hat{A}_{k+1}^{(\lambda,\ell)}$ such that $\mathfrak{v} \overset{i+\frac{1}{2}}{\sim} \mathfrak{t}$ and $\mathfrak{v}^{(i)} = \mathfrak{v}^{(i+1)}$. If $\mathfrak{s} \neq \mathfrak{v}$, then by item (1) of Proposition 2.5, and Lemma 5.9,

$$\begin{aligned}\sum_{\mathfrak{u} \overset{i+1}{\simeq} \mathfrak{t}} (c_{\mathfrak{u}}(i+1) - c_{\mathfrak{t}}(i))\sigma_{i+1}(\mathfrak{u},\mathfrak{t})f_{\mathfrak{u}}^{(k+1)} &= -f_{\mathfrak{t}}^{(k+1)}(L_ip_{i+\frac{1}{2}} + p_{i+\frac{1}{2}}p_{i+1} - p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} - 1)\\ &= f_{\mathfrak{t}}^{(k+1)} - c_{\mathfrak{t}}(i)f_{\mathfrak{t}}p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})f_{\mathfrak{v}}^{(k+1)}p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})c_i(\mathfrak{v})p_{i+1}(\mathfrak{v},\mathfrak{v})f_{\mathfrak{v}}p_{i+\frac{1}{2}},\end{aligned}$$

where

$$f_{\mathfrak{v}}^{(k+1)}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})^{-1}p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{v})f_{\mathfrak{t}}^{(k+1)}p_{i+\frac{1}{2}} \quad \text{and} \quad p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{v}) = p_{i+\frac{1}{2}}(\mathfrak{t},\mathfrak{t})p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{v}).$$

This gives the formula for $\sigma_{i+1}(\mathfrak{s},\mathfrak{t})$. Next,

$$f_{\mathfrak{t}}^{(k+1)}\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} = \sum_{\mathfrak{u} \overset{i+1}{\sim} \mathfrak{t}} \sigma_{i+1}(\mathfrak{u},\mathfrak{t})f_{\mathfrak{u}}^{(k+1)}p_{i+1}p_{i+\frac{1}{2}} = \sum_{\mathfrak{u} \overset{i+1}{\sim} \mathfrak{t}} \sigma_{i+1}(\mathfrak{u},\mathfrak{t})p_{i+1}(\mathfrak{v},\mathfrak{u})f_{\mathfrak{v}}^{(k+1)}p_{i+\frac{1}{2}}.$$

Since

$$f_{\mathfrak{v}}^{(k+1)}\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} = c_{\mathfrak{t}}(i)f_{\mathfrak{t}}^{(k+1)}p_{i+\frac{1}{2}} = c_{\mathfrak{t}}(i)p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{v})^{-1}p_{i+\frac{1}{2}}(\mathfrak{v},\mathfrak{t})f_{\mathfrak{v}}^{(k+1)}p_{i+\frac{1}{2}},$$

the formula for $\sigma_{i+1}(\mathfrak{v},\mathfrak{t})$ follows.

(2) Suppose that $\mathfrak{t}^{(i)} = \mathfrak{t}^{(i+1)}$ and $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+\frac{1}{2})}$, and let $\mathfrak{v} \in \hat{A}_{k+1}^{(\lambda,\ell)}$ such that $\mathfrak{v} \overset{i+1}{\sim} \mathfrak{t}$ and $\mathfrak{v}^{(i-\frac{1}{2})} = \mathfrak{v}^{(i+\frac{1}{2})}$. By item (1) of Proposition 2.5, and Lemma 5.9,

$$\sum_{\mathfrak{u} \overset{i+1}{\simeq} \mathfrak{t}} (c_{\mathfrak{u}}(i+1) - c_{\mathfrak{t}}(i))\sigma_{i+1}(\mathfrak{u},\mathfrak{t})f_{\mathfrak{u}}^{(k+1)} = f_{\mathfrak{t}}^{(k+1)},$$

which gives the formula for $\sigma_{i+1}(\mathfrak{s},\mathfrak{t})$ when $\mathfrak{s} \neq \mathfrak{v}$. Next,

$$f_{\mathfrak{t}}^{(k+1)}\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} = \sum_{\mathfrak{u} \overset{i+1}{\sim} \mathfrak{v}} \sigma_{i+1}(\mathfrak{u},\mathfrak{t})f_{\mathfrak{u}}^{(k+1)}p_{i+1}p_{i+\frac{1}{2}} = \sum_{\mathfrak{u} \overset{i+1}{\sim} \mathfrak{v}} \sigma_{i+1}(\mathfrak{u},\mathfrak{t})p_{i+1}(\mathfrak{v},\mathfrak{u})f_{\mathfrak{v}}^{(k+1)}p_{i+\frac{1}{2}}.$$

Since

$$f_{\mathfrak{t}}^{(k+1)}\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} = c_{\mathfrak{t}}(i)f_{\mathfrak{t}}^{(k+1)}p_{i+\frac{1}{2}} = 0 \qquad \text{and} \qquad f_{\mathfrak{v}}^{(k+1)}p_{i+\frac{1}{2}} \neq 0,$$

the formula for $\sigma_{i+1}(\mathfrak{v},\mathfrak{t})$ follows.

(3) Suppose that $\mathfrak{t}^{(i-\frac{1}{2})} \neq \mathfrak{t}^{(i+\frac{1}{2})}$ and $\mathfrak{t}^{(i)} \neq \mathfrak{t}^{(i+1)}$ and that $\mathfrak{t}\sigma_{i+1}$ does not exist. By item (1) of Proposition 2.5 and Lemma 5.9,

$$f_{\mathfrak{t}}^{(k+1)}\sigma_{i+1}L_{i+1} - f_{\mathfrak{t}}^{(k+1)}L_i\sigma_{i+\frac{1}{2}} = f_{\mathfrak{t}}^{(k+1)}.$$

Since

$$\mathfrak{u}\in\hat{A}^{(\lambda,\ell)}_{k+1} \text{ and } \mathfrak{u}\overset{i+1}{\simeq}\mathfrak{t} \qquad\qquad \text{implies that} \qquad\qquad \mathfrak{u}=\mathfrak{t},$$

it follows that $\sigma_{i+1}(\mathfrak{u},\mathfrak{t})=0$ if $\mathfrak{u}\neq\mathfrak{t}$, while

$$\left(c_{\mathfrak{t}}(i+1)-c_{\mathfrak{t}}(i)\right)\sigma_{i+1}(\mathfrak{t},\mathfrak{t})=1,$$

as required.
(4) Suppose that $\mathfrak{t}^{(i-\frac{1}{2})}\neq\mathfrak{t}^{(i+\frac{1}{2})}$ and $\mathfrak{t}^{(i)}\neq\mathfrak{t}^{(i+1)}$ and that $\mathfrak{t}\sigma_{i+1}$ exists. First observe that the hypotheses on $\mathfrak{t}$ imply that if $\mathfrak{u}\in\hat{A}^{(\lambda,\ell)}_{k+1}$ and $\mathfrak{u}\overset{i+1}{\simeq}\mathfrak{t}$, then $\mathfrak{u}\in\{\mathfrak{t},\mathfrak{t}\sigma_{i+1}\}$. Thus by Lemma (5.10),

$$f^{(k+1)}_{\mathfrak{t}}\sigma_{i+1}=\sigma_{i+1}(\mathfrak{t},\mathfrak{t})f^{(k+1)}_{\mathfrak{t}}+\sigma_{i+1}(\mathfrak{s},\mathfrak{t})f^{(k+1)}_{\mathfrak{s}}, \qquad\qquad \text{where } \mathfrak{s}=\mathfrak{t}\sigma_{i+1}.$$

We now determine $\sigma_{i+1}(\mathfrak{t},\mathfrak{t})$. Lemma 5.9 and the assumptions on $\mathfrak{t}$ imply that $f_{\mathfrak{t}}p_{i+1}=f_{\mathfrak{t}}p_{i+\frac{1}{2}}=0$. Thus item (1) of Proposition 2.5 gives

$$(c_{\mathfrak{t}}(i+1)-c_{\mathfrak{t}}(i))\sigma_{i+1}(\mathfrak{t},\mathfrak{t})=1,$$

as required.
Next, we show that if $\mathfrak{s}=\mathfrak{t}\sigma_{i+1}$ and $\mathfrak{t}\succ\mathfrak{s}$, then $\sigma(\mathfrak{s},\mathfrak{t})=1$. Now,

$$f^{(k+1)}_{\mathfrak{t}}\sigma_{i+1}=f^{(k+1)}_{\mathfrak{t}^{\lambda}}p_{\mathfrak{t}}\sigma_{i+1}+\sum_{\mathfrak{v}\succ\mathfrak{t}}r_{\mathfrak{v}}f^{(k+1)}_{\mathfrak{t}^{\lambda}}p_{\mathfrak{v}}\sigma_{i+1}, \qquad\qquad \text{where } r_{\mathfrak{v}}\in\kappa, \text{ for } \mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+1}.$$

It therefore suffices to show that

$$\sigma_{i+1}(\mathfrak{v},\mathfrak{s})=0, \qquad\qquad \text{if } \mathfrak{v}\in\hat{A}^{(\lambda,\ell)}_{k+1} \text{ and } \mathfrak{v}\succ\mathfrak{t}, \tag{5.16}$$

and that there exist $r_{\mathfrak{u}}\in\kappa$, for $\mathfrak{u}\in\hat{A}^{(\lambda,\ell)}_{k+1}$, such that

$$f^{(k+1)}_{\mathfrak{t}^{\lambda}}p_{\mathfrak{t}}\sigma_{i+1}=f^{(k+1)}_{\mathfrak{t}^{\lambda}}p_{\mathfrak{s}}+\sum_{\mathfrak{u}\succ\mathfrak{s}}r_{\mathfrak{u}}f^{(k+1)}_{\mathfrak{t}^{\lambda}}p_{\mathfrak{u}}. \tag{5.17}$$

To prove (5.16), suppose that $\mathfrak{v}\succ\mathfrak{t}$. Then

$$f^{(k+1)}_{\mathfrak{v}}\sigma_{i+1}=\sum_{\mathfrak{u}\overset{i+1}{\simeq}\mathfrak{v}}\sigma_{i+1}(\mathfrak{u},\mathfrak{v})f^{(k+1)}_{\mathfrak{u}}. \tag{5.18}$$

If $\sigma_{i+1}(\mathfrak{s},\mathfrak{v})\neq0$ in the expression (5.18), then $\mathfrak{v}\overset{i+1}{\simeq}\mathfrak{s}$, so $\mathfrak{v}\in\{\mathfrak{t},\mathfrak{s}\}$, which violates the assumption that $\mathfrak{v}\succ\mathfrak{t}\succ\mathfrak{s}$. Thus (5.16) holds.
Now we verify (5.17). There is no loss of generality in assuming that that $i=k$. There are two cases to consider here.
(a) If $\mathfrak{t}^{(k-\frac{1}{2})}=\mathfrak{t}^{(k)}$, let

$$(\mathfrak{t}^{(k-\frac{1}{2})},\mathfrak{t}^{(k)},\mathfrak{t}^{(k+\frac{1}{2})},\mathfrak{t}^{(k+1)})=(\mu,\mu,\nu,\lambda) \quad \text{and} \quad (\mathfrak{s}^{(k-\frac{1}{2})},\mathfrak{s}^{(k)},\mathfrak{s}^{(k+\frac{1}{2})},\mathfrak{s}^{(k+1)})=(\mu,\upsilon,\lambda,\lambda),$$

where

$$\lambda=\nu\cup\{(j,\lambda_j)\}, \qquad \mu=\nu\cup\{(j',\mu_{j'})\}, \qquad \text{and} \qquad \upsilon=\mu\cup\{(j,\upsilon_j)\}=\lambda\cup\{(j',\upsilon_{j'}).$$

If $\lambda\vdash k-\ell$, then

$$p_{\mathfrak{t}^{(k+\frac{1}{2})}\to\mathfrak{t}^{(k+1)}}=w_{a_j,k+1} \qquad \text{and} \qquad p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}}=w_{\ell-1,k}, \qquad \text{where} \qquad a_j=\ell+\sum_{r=1}^{j}\lambda_r,$$

and

$$p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}}=w_{\ell,k-1}p_{k+\frac{1}{2}}w_{k-1,a_{j'}}\sum_{r=0}^{\nu_{j'}}w_{a_{j'},a_{j'}-r}, \qquad \text{where} \qquad a_{j'}=\ell-1+\sum_{r=1}^{j'}\mu_r,$$

while

$$p_{\mathfrak{s}^{(k+\frac{1}{2})}\to\mathfrak{s}^{(k+1)}} = w_{\ell,k} \qquad \text{and} \qquad p_{\mathfrak{s}^{(k-\frac{1}{2})}\to\mathfrak{s}^{(k)}} = w_{\alpha_j,k-1}, \qquad \text{where} \qquad \alpha_j = \ell - 2 + \sum_{r=1}^{j} \upsilon_r,$$

and

$$p_{\mathfrak{s}^{(k)}\to\mathfrak{s}^{(k+\frac{1}{2})}} = w_{\ell-1,k-1} p_{k+\frac{1}{2}} w_{k-1,\alpha_{j'}} \sum_{r=0}^{\lambda_{j'}} w_{\alpha_{j'},\alpha_{j'}-r}, \qquad \text{where} \qquad \alpha_{j'} = \ell - 2 + \sum_{r=1}^{j'} \upsilon_r.$$

The relation $\sigma_{k+1} = s_k \sigma_{k+\frac{1}{2}}$ together with the braid relation, gives

$$\begin{aligned}
&f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{t}^{(k+\frac{1}{2})}\to\mathfrak{t}^{(k+1)}} p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}} p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}} \sigma_{k+1} \\
&\qquad = f_{\mathfrak{t}^\lambda}^{(k+1)} w_{a_j,k+1} w_{\ell,k-1} p_{k+\frac{1}{2}} w_{k-1,a_{j'}} \left(\sum_{r=0}^{\upsilon_{j'}} w_{a_{j'},a_{j'}-r} \right) w_{\ell-1,k+1} \sigma_{k+\frac{1}{2}} \\
&\qquad = f_{\mathfrak{t}^\lambda}^{(k+1)} w_{\ell-1,k+1} w_{a_j-1,k} w_{\ell-1,k-2} p_{k-\frac{1}{2}} w_{k-2,a_{j'}-1} \left(\sum_{r=0}^{\upsilon_{j'}} w_{a_{j'}-1,a_{j'}-r-1} \right) \sigma_{k+\frac{1}{2}} \\
&\qquad = f_{\mathfrak{t}^\lambda}^{(k+1)} w_{\ell,k+1} w_{a_j-1,k} w_{\ell-1,k-2} p_{k-\frac{1}{2}} w_{k-2,a_{j'}-1} \left(\sum_{r=0}^{\upsilon_{j'}} w_{a_{j'}-1,a_{j'}-r-1} \right) \sigma_{k+\frac{1}{2}} \\
&\qquad = f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{u}} \sigma_{k+\frac{1}{2}},
\end{aligned}$$

where $\mathfrak{u} \in \hat{A}_{k+1}^{(\lambda,\ell)}$ is given by

$$(\mathfrak{u}^{(k-1)}, \mathfrak{u}^{(k-\frac{1}{2})}, \mathfrak{u}^{(k)}, \mathfrak{u}^{(k+\frac{1}{2})}, \mathfrak{u}^{(k+1)}) = (\mu, \nu, \lambda, \lambda, \lambda)$$

and

$$p_{\mathfrak{u}^{(\frac{1}{2})}\to\mathfrak{u}^{(1)}} = p_{\mathfrak{u}^{(1)}\to\mathfrak{u}^{(1+\frac{1}{2})}} = \cdots = p_{\mathfrak{u}^{(k-\frac{3}{2})}\to\mathfrak{u}^{(k-1)}} = 1.$$

Since $\mathfrak{v} = \mathfrak{u}\sigma_{k+\frac{1}{2}}$ exists, and

$$(\mathfrak{v}^{(k-1)}, \mathfrak{v}^{(k-\frac{1}{2})}, \mathfrak{v}^{(k)}, \mathfrak{v}^{(k+\frac{1}{2})}, \mathfrak{v}^{(k+1)}) = (\mu, \mu, \upsilon, \lambda, \lambda),$$

it follows that $\mathfrak{u} \succ \mathfrak{v}$. By (5.12), there exist $r_{\mathfrak{a}} \in \kappa$, for $\mathfrak{a} \in \hat{A}_{k+1}^{(\lambda,\ell)}$, such that

$$f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{t}^{(k+\frac{1}{2})}\to\mathfrak{t}^{(k+1)}} p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}} p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}} \sigma_{k+1} = f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{v}} + \sum_{\mathfrak{a}\succ\mathfrak{v}} r_{\mathfrak{a}} f_{\mathfrak{t}^\lambda}^{(k+1)} p_{\mathfrak{a}}.$$

Multiplying both sides of the last expression by

$$p_{\mathfrak{s}^{(k-1)}\to\mathfrak{s}^{(k-\frac{1}{2})}} p_{\mathfrak{s}^{(k-\frac{3}{2})}\to\mathfrak{s}^{(k-1)}} \cdots p_{\mathfrak{s}^{(\frac{1}{2})}\to\mathfrak{s}^{(1)}} = p_{\mathfrak{t}^{(k-1)}\to\mathfrak{t}^{(k-\frac{1}{2})}} p_{\mathfrak{t}^{(k-\frac{3}{2})}\to\mathfrak{t}^{(k-1)}} \cdots p_{\mathfrak{t}^{(\frac{1}{2})}\to\mathfrak{t}^{(1)}}$$

on the right verifies (5.17) in this instance.
(b) If $\mathfrak{t}^{(k)} = \mathfrak{t}^{(k+\frac{1}{2})}$, let $\mathfrak{s} = \mathfrak{t}\sigma_{k+1}$,

$$(\mathfrak{s}^{(k-\frac{1}{2})}, \mathfrak{s}^{(k)}, \mathfrak{s}^{(k+\frac{1}{2})}, \mathfrak{s}^{(k+1)}) = (\mu, \nu, \nu, \lambda) \qquad \text{and} \qquad (\mathfrak{t}^{(k-\frac{1}{2})}, \mathfrak{t}^{(k)}, \mathfrak{t}^{(k+\frac{1}{2})}, \mathfrak{t}^{(k+1)}) = (\mu, \upsilon, \upsilon, \lambda)$$

where

$$\lambda = \nu \cup \{(j, \lambda_j)\}, \qquad \nu = \mu \cup \{(j', \lambda_{j'}\}, \qquad \text{and} \qquad \lambda = \upsilon \cup \{(j', \lambda_{j'})\}, \qquad \upsilon = \mu \cup \{(j, \lambda_j\}.$$

and $j' < j$. If

$$a_j = \ell + \sum_{r=1}^{j} \lambda_r, \qquad \text{and} \qquad a_{j'} = \ell + \sum_{r=1}^{j'} \lambda_r,$$

then

$$p_{\mathfrak{s}^{(k+\frac{1}{2})}\to\mathfrak{s}^{(k+1)}}p_{\mathfrak{s}^{(k)}\to\mathfrak{s}^{(k+\frac{1}{2})}}p_{\mathfrak{s}^{(k-\frac{1}{2})}\to\mathfrak{s}^{(k)}} = w_{a_{j'},k}w_{a_j,k-1}$$

and

$$p_{\mathfrak{t}^{(k+\frac{1}{2})}\to\mathfrak{t}^{(k+1)}}p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}}p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}} = w_{a_{j'},k}w_{a_j-1,k-1}.$$

The braid relation and $s_{k-1}\sigma_{k+1} = \sigma_{k+\frac{1}{2}}$ give

$$\begin{aligned}p_{\mathfrak{t}^{(k+\frac{1}{2})}\to\mathfrak{s}^{(k+1)}}p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}}p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}}\sigma_{k+1} &= w_{a_j,k}w_{a_{j'}-1,k-1}\sigma_{k+1}\\ &= p_{\mathfrak{s}^{(k+\frac{1}{2})}\to\mathfrak{s}^{(k+1)}}p_{\mathfrak{s}^{(k)}\to\mathfrak{s}^{(k+\frac{1}{2})}}p_{\mathfrak{s}^{(k-\frac{1}{2})}\to\mathfrak{s}^{(k)}}\sigma_{k+\frac{1}{2}}.\end{aligned}$$

By (5.12), there exist $r_{\mathfrak{a}} \in \kappa$, for $\mathfrak{a} \in \hat{A}_{k+1}^{(\lambda,\ell)}$, such that

$$f_{\mathfrak{t}^\lambda}^{(k+1)}p_{\mathfrak{t}^{(k+\frac{1}{2})}\to\mathfrak{t}^{(k+1)}}p_{\mathfrak{t}^{(k)}\to\mathfrak{t}^{(k+\frac{1}{2})}}p_{\mathfrak{t}^{(k-\frac{1}{2})}\to\mathfrak{t}^{(k)}}\sigma_{k+1} = f_{\mathfrak{t}^\lambda}^{(k+1)}p_{\mathfrak{v}} + \sum_{\mathfrak{a}\succ\mathfrak{v}} r_{\mathfrak{a}}f_{\mathfrak{t}^\lambda}^{(k+1)}p_{\mathfrak{a}}, \tag{5.19}$$

where $\mathfrak{v} \in \hat{A}_{k+1}^{(\lambda,\ell)}$ is given by

$$p_{\mathfrak{v}} = p_{\mathfrak{s}^{(k+\frac{1}{2})}\to\mathfrak{s}^{(k+1)}}p_{\mathfrak{s}^{(k)}\to\mathfrak{s}^{(k+\frac{1}{2})}}p_{\mathfrak{s}^{(k-\frac{1}{2})}\to\mathfrak{s}^{(k)}}.$$

Multiplying both sides of the expression (5.19) by

$$p_{\mathfrak{s}^{(k-1)}\to\mathfrak{s}^{(k-\frac{1}{2})}}p_{\mathfrak{s}^{(k-\frac{3}{2})}\to\mathfrak{s}^{(k-1)}}\cdots p_{\mathfrak{s}^{(\frac{1}{2})}\to\mathfrak{s}^{(1)}} = p_{\mathfrak{t}^{(k-1)}\to\mathfrak{t}^{(k-\frac{1}{2})}}p_{\mathfrak{t}^{(k-\frac{3}{2})}\to\mathfrak{t}^{(k-1)}}\cdots p_{\mathfrak{t}^{(\frac{1}{2})}\to\mathfrak{t}^{(1)}}$$

on the right completes the proof (5.17) and establishes that $\sigma_{i+1}(\mathfrak{s},\mathfrak{t}) = 1$ if $\mathfrak{s} = \mathfrak{t}\sigma_{i+1}$ and $\mathfrak{t} \succ \mathfrak{s}$. Finally, using the above calculations together with the fact that $\sigma_{i+1}^2 = 1$, we can verify the formula for $\sigma_{i+1}(\mathfrak{s},\mathfrak{t})$ when $\mathfrak{s} = \mathfrak{t}\sigma_{i+\frac{1}{2}}$ and $\mathfrak{s} \succ \mathfrak{t}$ by observing that $c_{\mathfrak{s}}(i) = c_{\mathfrak{t}}(i+1)$ and $c_{\mathfrak{s}}(i+1) = c_{\mathfrak{t}}(i)$. $\square$

6. Central Element Recursions

In this section we obtain partition algebra analogues of the central element recursions obtained by Nazarov [Na] for the Brauer algebras and Beliakova and Blanchet [BB] for the BMW algebras, and explain the relationship between these central element recursions and the seminormal representations of the partition algebras.

We renormalise the Jucys–Murphy elements (2.3) and (2.5) by defining elements

$$x_{i+\frac{1}{2}} = -\tfrac{z}{2} + L_{i+\frac{1}{2}} \qquad \text{and} \qquad x_{i+1} = -\tfrac{z}{2} + L_{i+1}, \qquad \text{for } i = 0, 1, \ldots.$$

Then, for $i = 1, 2, \ldots$, we have the relations

$$x_{i+\frac{1}{2}} = -x_i p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}x_i - (\tfrac{z}{2} + x_{i-\frac{1}{2}})p_{i+\frac{1}{2}} + s_i x_{i-\frac{1}{2}} s_i + \sigma_{i+\frac{1}{2}}, \tag{6.1}$$

$$x_{i+1} = -\tfrac{z}{2}p_{i+\frac{1}{2}} - s_i x_i p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}x_i s_i + p_{i+\frac{1}{2}}x_i p_{i+1} p_{i+\frac{1}{2}} + s_i x_{i-1} s_i + \sigma_{i+1}. \tag{6.2}$$

Following §2 of [Na], for $i = 0, 1, \ldots$, and $j = 0, 1, \ldots$, define central elements

$$x_{i+\frac{1}{2}}^{(j)} \in \mathcal{A}_i(z) \qquad \text{and} \qquad x_{i+1}^{(j)} \in \mathcal{A}_{i+\frac{1}{2}}(z).$$

by

$$x_{i+\frac{1}{2}}^{(j)}p_{i+1} = p_{i+1}(x_{i+\frac{1}{2}})^j p_{i+1} \qquad \text{and} \qquad x_{i+1}^{(j)}p_{i+\frac{3}{2}} = p_{i+\frac{3}{2}}(x_{i+1})^j p_{i+\frac{3}{2}}. \tag{6.3}$$

Let u denote a formal variable, and define the central elements $W_{i+\frac{1}{2}}(u) \in \mathcal{A}_i(z)[[u^{-1}]]$, and $W_{i+1}(u) \in \mathcal{A}_{i+\frac{1}{2}}(z)[[u^{-1}]]$, for $i = 0, 1, \ldots$, by

$$W_{i+\frac{1}{2}}(u) = u^{-1}\sum_{j\geqslant 0} x_{i+\frac{1}{2}}^{(j)}u^{-j} \qquad \text{and} \qquad W_{i+1}(u) = \sum_{j\geqslant 0} x_{i+1}^{(j)}u^{-j}.$$

For $i = 0, 1, \dots,$ we will work with the formal expressions

$$W_{i+\frac{1}{2}}(u)p_{i+1} = p_{i+1}\frac{1}{u - x_{i+\frac{1}{2}}}p_{i+1} \qquad \text{and} \qquad W_{i+1}(u)p_{i+\frac{3}{2}} = p_{i+\frac{3}{2}}\frac{u}{u - x_{i+1}}p_{i+\frac{3}{2}}.$$

The next statement is the partition algbebra analogue of Proposition 4.2 of [Na] and (12) of [BB].

Proposition 6.1. *If* $i = 1, 2, \dots,$ *then*

$$\frac{W_{i+\frac{1}{2}}(u) + (\frac{z}{2} - u - 1)}{W_{i-\frac{1}{2}}(u) + (\frac{z}{2} - u - 1)} = \frac{(u + x_i)^2 - 1}{(u - x_{i-\frac{1}{2}})^2 - 1}\,\frac{(u - x_{i-\frac{1}{2}})^2}{(u + x_i)^2}, \tag{6.4}$$

and

$$\frac{W_{i+1}(u)}{W_i(u)} = \frac{(u + x_{i+\frac{1}{2}})^2 - 1}{(u - x_i)^2 - 1}\,\frac{(u - x_i)^2}{(u + x_{i+\frac{1}{2}})^2}. \tag{6.5}$$

Proof. We first prove (6.4). From the relation (6.1),

$$\begin{aligned} s_i(u - x_{i+\frac{1}{2}}) &= \tfrac{z}{2}p_{i+\frac{1}{2}} + s_i x_i p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}}x_i + x_{i-\frac{1}{2}}p_{i+\frac{1}{2}} + (u - x_{i-\frac{1}{2}})s_i - \sigma_{i+1} \\ &= \sigma_{i+1}p_i p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}}x_i + x_{i-\frac{1}{2}}p_{i+\frac{1}{2}} + (u - x_{i-\frac{1}{2}})s_i - \sigma_{i+1}, \end{aligned}$$

and

$$\begin{aligned} \frac{1}{u - x_{i-\frac{1}{2}}}s_i = {}& \frac{1}{u - x_{i-\frac{1}{2}}}\sigma_{i+1}p_i p_{i+\frac{1}{2}}\frac{1}{u - x_{i+\frac{1}{2}}} + \frac{1}{u - x_{i-\frac{1}{2}}}p_{i+\frac{1}{2}}x_i\frac{1}{u - x_{i+\frac{1}{2}}} \\ & - p_{i+\frac{1}{2}}\frac{1}{u - x_{i+\frac{1}{2}}} + \frac{u}{u - x_{i-\frac{1}{2}}}p_{i+\frac{1}{2}}\frac{1}{u - x_{i+\frac{1}{2}}} + s_i\frac{1}{u - x_{i+\frac{1}{2}}} \\ & - \sigma_{i+1}\frac{1}{(u - x_{i+\frac{1}{2}})(u - x_{i-\frac{1}{2}})}. \end{aligned}$$

Using the fact that $p_{i+\frac{1}{2}}(u - x_{i+\frac{1}{2}}) = p_{i+\frac{1}{2}}(u + x_i)$, the previous expression gives

$$\begin{aligned} \frac{1}{u - x_{i-\frac{1}{2}}}s_i = {}& \frac{1}{u - x_{i-\frac{1}{2}}}\sigma_{i+1}p_i p_{i+\frac{1}{2}}\frac{1}{u - x_{i+\frac{1}{2}}} + \frac{1}{u - x_{i-\frac{1}{2}}}p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}\frac{1}{u + x_i} \\ & + s_i\frac{1}{u - x_{i+\frac{1}{2}}} - \sigma_{i+1}\frac{1}{(u - x_{i+\frac{1}{2}})(u - x_{i-\frac{1}{2}})}. \end{aligned}$$

Since σ_{i+1} commutes with $x_{i-\frac{1}{2}}$, multiplying both sides of the last expression by σ_{i+1} on the left, we obtain

$$\begin{aligned} \frac{1}{u - x_{i-\frac{1}{2}}}\sigma_{i+\frac{1}{2}} = {}& \frac{1}{u - x_{i-\frac{1}{2}}}p_i p_{i+\frac{1}{2}}\frac{1}{u - x_{i+\frac{1}{2}}} + \frac{1}{u - x_{i-\frac{1}{2}}}p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}\frac{1}{u + x_i} \\ & + \sigma_{i+\frac{1}{2}}\frac{1}{u - x_{i+\frac{1}{2}}} - \frac{1}{(u - x_{i+\frac{1}{2}})(u - x_{i-\frac{1}{2}})}. \end{aligned} \tag{6.6}$$

Multiplying each side of (6.6) by p_{i+1} on the right and on the left,

$$\begin{aligned} p_{i+1}\sigma_{i+\frac{1}{2}}\frac{1}{u - x_{i+\frac{1}{2}}}p_{i+1} = {}& \frac{1}{u - x_{i-\frac{1}{2}}}p_{i+1}\sigma_{i+\frac{1}{2}}p_{i+1} - \frac{1}{u - x_{i-\frac{1}{2}}}p_i\frac{1}{u + x_i}p_{i+1} \\ & + \frac{1}{u + x_i}p_{i+1} - \frac{1}{u - x_{i-\frac{1}{2}}}p_{i+1} + \frac{1}{u - x_{i-\frac{1}{2}}}W_{i+\frac{1}{2}}(u)p_{i+1}. \end{aligned} \tag{6.7}$$

Using (15) of Proposition 2.4, we obtain $p_{i+1}\sigma_{i+\frac{1}{2}}p_{i+1} = (\frac{z}{2} - x_{i-\frac{1}{2}})p_{i+1}$ which, substituted into (6.7), gives

$$\begin{aligned} p_{i+1}\sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i+\frac{1}{2}}}p_{i+1} &= (\tfrac{z}{2} - u - 1)\frac{1}{u-x_{i-\frac{1}{2}}}p_{i+1} + p_{i+1} - \frac{1}{u-x_{i-\frac{1}{2}}}p_i\frac{1}{u+x_i}p_{i+1} \\ &\quad + \frac{1}{u+x_i}p_{i+1} - \frac{1}{u-x_{i-\frac{1}{2}}}p_{i+1} + \frac{1}{u-x_{i-\frac{1}{2}}}W_{i+\frac{1}{2}}(u)p_{i+1}. \end{aligned} \tag{6.8}$$

Multiplying each side of (6.6) by p_i on the left, and applying the algebra anti-involution $*$ to the result,

$$\begin{aligned} \sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i-\frac{1}{2}}}p_i &= \frac{1}{u+x_i}p_{i+\frac{1}{2}}p_iW_{i-\frac{1}{2}}(u) + \frac{1}{u-x_{i-\frac{1}{2}}}p_{i+\frac{1}{2}}p_i - \frac{1}{u+x_i}p_{i+\frac{1}{2}}p_i \\ &\quad + \frac{1}{u-x_{i+\frac{1}{2}}}\sigma_{i+\frac{1}{2}}p_i - \frac{1}{(u+x_i)(u-x_{i+\frac{1}{2}})}p_i. \end{aligned}$$

Making the substitution $\sigma_{i+\frac{1}{2}}p_i = (\frac{z}{2} + x_i)p_{i+\frac{1}{2}}p_i$ in the last expression gives

$$\begin{aligned} \sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i-\frac{1}{2}}}p_i &= \frac{1}{u+x_i}p_{i+\frac{1}{2}}p_iW_{i-\frac{1}{2}}(u) + \frac{1}{u-x_{i-\frac{1}{2}}}p_{i+\frac{1}{2}}p_i + p_{i+\frac{1}{2}}p_i \\ &\quad + (\tfrac{z}{2} - u - 1)\frac{1}{u+x_i}p_{i+\frac{1}{2}}p_i - \frac{1}{(u+x_i)(u-x_{i+\frac{1}{2}})}p_i. \end{aligned}$$

Therefore,

$$\begin{aligned} p_{i+1}\sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i-\frac{1}{2}}}p_ip_{i+\frac{1}{2}}\frac{1}{u+x_i}p_{i+1} &= \frac{1}{(u+x_i)^2}W_{i-\frac{1}{2}}(u)p_{i+1} + \frac{1}{u+x_i}p_{i+1} \\ + (\tfrac{z}{2} - u - 1)\frac{1}{(u+x_i)^2}p_{i+1} &+ \frac{1}{(u-x_{i-\frac{1}{2}})(u+x_i)}p_{i+1} \\ - \frac{1}{u-x_{i-\frac{1}{2}}}p_i\frac{1}{(u+x_i)^2}p_{i+1}&. \end{aligned} \tag{6.9}$$

Now,

$$\begin{aligned} p_{i+1}\sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i-\frac{1}{2}}}\sigma_{i+\frac{1}{2}}p_{i+1} &= p_{i+1}\sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i-\frac{1}{2}}}p_ip_{i+\frac{1}{2}}\frac{1}{u-x_{i+\frac{1}{2}}}p_{i+1} \\ &\quad + \frac{1}{u-x_{i-\frac{1}{2}}}p_{i+1} - \frac{1}{u+x_i}p_{i+1} + W_{i+\frac{1}{2}}(u)p_{i+1} \\ &\quad - p_{i+1}\sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i+\frac{1}{2}}}p_{i+1}\frac{1}{u-x_{i-\frac{1}{2}}}. \end{aligned} \tag{6.10}$$

Using the relation

$$p_{i+1}\sigma_{i+\frac{1}{2}}\frac{1}{u-x_{i-\frac{1}{2}}}\sigma_{i+\frac{1}{2}}p_{i+1} = s_ip_i\sigma_{i+1}\frac{1}{u-x_{i-\frac{1}{2}}}\sigma_{i+1}p_is_i = W_{i-\frac{1}{2}}(u)p_{i+1},$$

and substituting (6.8) and (6.9) into (6.10), we obtain

$$\begin{aligned} W_{i-\frac{1}{2}}(u)p_{i+1} &= \frac{1}{(u+x_i)^2}W_{i-\frac{1}{2}}(u)p_{i+1} + (\tfrac{z}{2} - u - 1)\frac{1}{(u+x_i)^2}p_{i+1} + W_{i+\frac{1}{2}}(u)p_{i+1} \\ &\quad - \frac{1}{(u-x_{i-\frac{1}{2}})^2}p_{i+1} - (\tfrac{z}{2} - u - 1)\frac{1}{(u-x_{i-\frac{1}{2}})^2}p_{i+1} \end{aligned}$$

or

$$\frac{(u+x_i)^2-1}{(u+x_i)^2}p_{i+1}W_{i-\frac{1}{2}}(u)-(\tfrac{z}{2}-u-1)\frac{1}{(u+x_i)^2}p_{i+1}$$
$$=\frac{(u-x_{i-\frac{1}{2}})^2-1}{(u-x_{i-\frac{1}{2}})^2}p_{i+1}W_{i+\frac{1}{2}}(u)-(\tfrac{z}{2}-u-1)\frac{1}{(u-x_{i-\frac{1}{2}})^2}p_{i+1},$$

from which the relation (6.4) now follows.
For the proof of(6.5), use the the relation (6.2) to write

$$(u-x_{i+1})s_i=\tfrac{z}{2}p_{i+\frac{1}{2}}+s_ix_ip_{i+\frac{1}{2}}+p_{i+\frac{1}{2}}x_i-p_{i+\frac{1}{2}}x_ip_{i+1}p_{i+\frac{1}{2}}+s_i(u-x_i)-\sigma_{i+\frac{1}{2}}. \tag{6.11}$$

Since $(x_{i+1})^jp_{i+\frac{1}{2}}=p_{i+\frac{1}{2}}(x_i)^jp_{i+1}p_{i+\frac{1}{2}}$ for $j=0,1,\dots$, we obtain

$$\frac{1}{u-x_{i+1}}p_{i+\frac{1}{2}}=p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+1}p_{i+\frac{1}{2}},$$

which together with (6.11), gives

$$\begin{aligned}s_i\frac{1}{u-x_i}&=\frac{z}{2}\frac{1}{u-x_{i+1}}p_{i+\frac{1}{2}}\frac{1}{u-x_i}+\frac{1}{u-x_{i+1}}s_ix_ip_{i+\frac{1}{2}}\frac{1}{u-x_i}-p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+1}p_{i+\frac{1}{2}}\\&\quad+p_{i+\frac{1}{2}}\frac{1}{u-x_i}+\frac{1}{u-x_{i+1}}s_i-\sigma_{i+\frac{1}{2}}\frac{1}{(u-x_i)(u-x_{i+1})}\\&=\frac{1}{u-x_{i+1}}\sigma_{i+\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}}\frac{1}{u-x_i}-p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+1}p_{i+\frac{1}{2}}+p_{i+\frac{1}{2}}\frac{1}{u-x_i}\\&\quad+\frac{1}{u-x_{i+1}}s_i-\sigma_{i+\frac{1}{2}}\frac{1}{(u-x_i)(u-x_{i+1})}.\end{aligned}$$

Multiplying both sides of the above expression by $\sigma_{i+\frac{1}{2}}$ on the left,

$$\begin{aligned}\sigma_{i+1}\frac{1}{u-x_i}&=\frac{1}{u-x_{i+1}}p_{i+1}p_{i+\frac{1}{2}}\frac{1}{u-x_i}-p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+1}p_{i+\frac{1}{2}}+p_{i+\frac{1}{2}}\frac{1}{u-x_i}\\&\quad+\frac{1}{u-x_{i+1}}\sigma_{i+1}-\frac{1}{(u-x_i)(u-x_{i+1})},\end{aligned}\tag{6.12}$$

and multiplying both sides of the expression (6.12) by $p_{i+\frac{1}{2}}$ on the right, and then applying the anti-involution $*$ to the result,

$$\begin{aligned}p_{i+\frac{1}{2}}\frac{1}{u-x_i}\sigma_{i+1}&=\frac{W_i(u)}{u}p_{i+\frac{1}{2}}p_{i+1}\frac{1}{u-x_{i+1}}+\frac{W_i(u)}{u}p_{i+\frac{1}{2}}\\&\quad-p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+1}p_{i+\frac{1}{2}}\frac{1}{u-x_i}.\end{aligned}\tag{6.13}$$

Multiplying both sides of the expression (6.12) by $p_{i+\frac{3}{2}}$ on the right and on the left, and then using (17) of Proposition 2.4 together with the fact that $p_{i+1}(u+x_{i+\frac{1}{2}})=p_{i+1}(u-x_{i+1})$, we obtain

$$\begin{aligned}p_{i+\frac{3}{2}}\frac{1}{u-x_{i+1}}\sigma_{i+1}p_{i+\frac{3}{2}}&=p_{i+\frac{1}{2}}p_{i+\frac{3}{2}}\frac{W_i(u)}{u}-\frac{1}{u+x_{i+\frac{1}{2}}}p_{i+\frac{1}{2}}p_{i+\frac{3}{2}}\frac{1}{u-x_i}\\&\quad+\frac{1}{u-x_i}p_{i+\frac{3}{2}}\frac{W_{i+1}(u)}{u}.\end{aligned}\tag{6.14}$$

Multiplying (6.12) on the right by σ_{i+1} and substituting for the term appearing on the left hand side of (6.13),

$$\begin{aligned}\sigma_{i+1}\frac{1}{u-x_i}\sigma_{i+1} &= \frac{W_i(u)}{u}\frac{1}{u+x_{i+\frac{1}{2}}}p_{i+1}\frac{1}{u+x_{i+\frac{1}{2}}} + \frac{W_i(u)}{u}\frac{1}{u+x_{i+\frac{1}{2}}}p_{i+1}p_{i+\frac{1}{2}} \\ &\quad - \frac{1}{(u-x_i)(u+x_{i+\frac{1}{2}})}p_{i+1}p_{i+\frac{1}{2}}\frac{1}{u-x_i} - p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+1}p_{i+\frac{1}{2}} \\ &\quad + \frac{W_i(u)}{u}p_{i+\frac{1}{2}}p_{i+1}\frac{1}{u+x_{i+\frac{1}{2}}} + \frac{W_i(u)}{u}p_{i+\frac{1}{2}} \\ &\quad - p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+1}p_{i+\frac{1}{2}}\frac{1}{u-x_i} + \frac{1}{u-x_{i+1}} - \frac{1}{(u-x_i)(u-x_{i+1})}\sigma_{i+1}. \end{aligned} \tag{6.15}$$

Since

$$\begin{aligned} p_{i+\frac{3}{2}}\sigma_{i+1}\frac{1}{u-x_i}\sigma_{i+1}p_{i+\frac{3}{2}} &= p_{i+\frac{3}{2}}\sigma_{i+1}\sigma_{i+\frac{3}{2}}\frac{1}{u-x_i}\sigma_{i+\frac{3}{2}}\sigma_{i+1}p_{i+\frac{3}{2}} \\ &= \sigma_{i+1}\sigma_{i+\frac{3}{2}}p_{i+\frac{1}{2}}\frac{1}{u-x_i}p_{i+\frac{1}{2}}\sigma_{i+\frac{3}{2}}\sigma_{i+1} \\ &= \sigma_{i+1}\sigma_{i+\frac{3}{2}}p_{i+\frac{1}{2}}\sigma_{i+\frac{3}{2}}\sigma_{i+1}\frac{W_i(u)}{u} = p_{i+\frac{3}{2}}\frac{W_i(u)}{u}, \end{aligned}$$

multiplying both sides of (6.15) by $p_{i+\frac{3}{2}}$ on the left and on the right, and substituting for the term appearing on the left hand side of (6.14) gives

$$\begin{aligned} p_{i+\frac{3}{2}}\frac{W_i(u)}{u} = p_{i+\frac{3}{2}}\Bigg(&\frac{W_i(u)}{u}\frac{1}{(u+x_{i+\frac{1}{2}})^2} + \frac{W_i(u)}{u}\frac{1}{u+x_{i+\frac{1}{2}}}p_{i+\frac{1}{2}} \\ &- \frac{1}{(u-x_i)(u+x_{i+\frac{1}{2}})}p_{i+\frac{1}{2}}\frac{1}{u-x_i} - \frac{W_i(u)}{u}p_{i+\frac{1}{2}} + \frac{W_i(u)}{u}p_{i+\frac{1}{2}}\frac{1}{u+x_{i+\frac{1}{2}}} \\ &+ \frac{W_i(u)}{u}p_{i+\frac{1}{2}} - \frac{W_i(u)}{u}p_{i+\frac{1}{2}}\frac{1}{u-x_i} + \frac{W_{i+1}(u)}{u} - \frac{1}{u-x_i}\frac{W_i(u)}{u}p_{i+\frac{1}{2}} \\ &+ \frac{1}{(u-x_i)(u+x_{i+\frac{1}{2}})}p_{i+\frac{1}{2}}\frac{1}{u-x_i} - \frac{1}{(u-x_i)^2}\frac{W_{i+1}(u)}{u}\Bigg). \end{aligned}$$

Since $(u-x_i)p_{i+\frac{1}{2}} = (u+x_{i+\frac{1}{2}})p_{i+\frac{1}{2}}$, we obtain

$$\frac{W_i(u)}{u} - \frac{W_i(u)}{u}\frac{1}{(u+x_{i+\frac{1}{2}})^2} = \frac{W_{i+1}(u)}{u} - \frac{W_{i+1}(u)}{u}\frac{1}{(u-x_i)^2},$$

and the statement (6.5) follows. □

As an application of the recursions (6.4) and (6.5), the seminormal matrix entries of the contractions p_i and $p_{i+\frac{1}{2}}$ can be computed independently of any formula for the dimensions of the irreducible representations of the symmetric group.

Determine a series $Q_{i+\frac{1}{2}}(u) \in A_i(z)[[u^{-1}]]$ by the recursion (6.4) and

$$Q_{i+\frac{1}{2}}(u) = Z_{i+\frac{1}{2}}(u) + (\tfrac{z}{2} - u - 1), \qquad \text{and} \qquad Q_{\frac{1}{2}}(u) = -\frac{(u+1+\frac{z}{2})(u-\frac{z}{2})}{(u+\frac{z}{2})}, \tag{6.16}$$

and a series $Q_{i+1}(u) \in A_{i+\frac{1}{2}}(z)[[u^{-1}]]$ by the recursion (6.5), and

$$Q_{i+1}(u) = \frac{W_{i+1}(u)}{u}, \qquad \text{and} \qquad Q_1(u) = \frac{(u+1-\frac{z}{2})}{(u+\frac{z}{2})(u-\frac{z}{2})}. \tag{6.17}$$

If $i = 0, 1, \dots,$ and $\mu \in \hat{A}_i$, denote by

$Q_{i+\frac{1}{2}}(u, \mu)$ the scalar by which $Q_{i+\frac{1}{2}}(u)$ acts on the $A_i(z)[[u^{-1}]]$–module $A_i^{(\mu,m)}$.

Similarly, if $\mu \in \hat{A}_{i+\frac{1}{2}}$, denote by

$Q_{i+1}(u, \mu)$ the scalar by which $Q_{i+1}(u)$ acts on the $A_{i+\frac{1}{2}}(z)[[u^{-1}]]$–module $A_{i+\frac{1}{2}}^{(\mu,m)}$.

Given Proposition 3.4, the diagonal entries of the contractions p_i and $p_{i+\frac{1}{2}}$ in the seminormal matrix representation of $A_{k+\frac{1}{2}}(z)$ are determined by the recursions (6.16) and (6.17) as follows. If $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$ and $\mathfrak{t} = (\mu^{(0)}, \mu^{(\frac{1}{2})}, \dots, \mu^{(k+\frac{1}{2})}) \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$, define

$$x_{\mathfrak{t}}(i) = \begin{cases} \frac{z}{2} - |\mu^{(i)}|, & \text{if } \mu^{(i)} = \mu^{(i-\frac{1}{2})} \\ -\frac{z}{2} + c(b), & \text{if } \mu^{(i)} = \mu^{(i-\frac{1}{2})} \cup \{b\}, \end{cases} \qquad \text{for } i = 0, 1, \dots, k,$$

and

$$x_{\mathfrak{t}}(i - \tfrac{1}{2}) = \begin{cases} -\frac{z}{2} + |\mu^{(i-1)}|, & \text{if } \mu^{(i-1)} = \mu^{(i-\frac{1}{2})}, \\ \frac{z}{2} - c(b), & \text{if } \mu^{(i+\frac{1}{2})} = \mu^{(i)} \setminus \{b\}, \end{cases} \qquad \text{for } i = 1, \dots, k+1$$

which are the eigenvalues of the Jucys–Murphy elements under the normalisation (6.1) and (6.2). The next statement is a counterpart to (3.6) of [Na] and Lemma 7.4 of [BB].

Proposition 6.2. *Let $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$. The following statements hold:*

(1) If $\mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ satisfies $\mathfrak{t}^{(i-\frac{1}{2})} = \mathfrak{t}^{(i+\frac{1}{2})} = \mu$, then

$$p_{i+\frac{1}{2}}(\mathfrak{t}, \mathfrak{t}) = \operatorname{Res}_{u=x_{\mathfrak{t}}(i)} Q_i(u, \mu), \qquad \textit{for } i = 1, \dots, k.$$

(2) If $\mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ satisfies $\mathfrak{t}^{(i-1)} = \mathfrak{t}^{(i)} = \mu$, then

$$p_i(\mathfrak{t}, \mathfrak{t}) = \operatorname{Res}_{u=x_{\mathfrak{t}}(i-\frac{1}{2})} Q_{i-\frac{1}{2}}(u, \mu), \qquad \textit{for } i = 1, \dots, k.$$

Proof. (1) If $(\lambda, \ell) \in \hat{A}_{k+\frac{1}{2}}$ and $\mathfrak{t} \in \hat{A}_{k+\frac{1}{2}}^{(\lambda,\ell)}$ are as in the statement of the proposition, then

$$\begin{aligned} f_{\mathfrak{t}} \frac{W_i(u)}{u} p_{i+\frac{1}{2}} &= f_{\mathfrak{t}} p_{i+\frac{1}{2}} \frac{1}{u - x_i} p_{i+\frac{1}{2}} = \sum_{\mathfrak{s} \overset{i+\frac{1}{2}}{\sim} \mathfrak{t}} \frac{p_{i+\frac{1}{2}}(\mathfrak{s}, \mathfrak{t})}{u - x_{\mathfrak{s}}(i)} f_{\mathfrak{s}} p_{i+\frac{1}{2}} = \sum_{\substack{\mathfrak{s} \overset{i+\frac{1}{2}}{\sim} \mathfrak{t}, \\ \mathfrak{v} \overset{i+\frac{1}{2}}{\sim} \mathfrak{s}}} \frac{p_{i+\frac{1}{2}}(\mathfrak{v}, \mathfrak{s}) p_{i+\frac{1}{2}}(\mathfrak{s}, \mathfrak{t})}{u - x_{\mathfrak{s}}(i)} f_{\mathfrak{v}} \\ &= \sum_{\mathfrak{s} \overset{i+\frac{1}{2}}{\sim} \mathfrak{t}} \sum_{\mathfrak{v} \overset{i+\frac{1}{2}}{\sim} \mathfrak{s}} \frac{p_{i+\frac{1}{2}}(\mathfrak{s}, \mathfrak{s}) p_{i+\frac{1}{2}}(\mathfrak{v}, \mathfrak{t})}{u - x_{\mathfrak{s}}(i)} f_{\mathfrak{v}} = \sum_{\mathfrak{s} \overset{i+\frac{1}{2}}{\sim} \mathfrak{t}} \frac{p_{i+\frac{1}{2}}(\mathfrak{s}, \mathfrak{s})}{u - x_{\mathfrak{s}}(i)} f_{\mathfrak{t}} p_{i+\frac{1}{2}}, \end{aligned}$$

together with the fact that $p_{i+\frac{1}{2}}(\mathfrak{t}, \mathfrak{t}) \neq 0$, gives the required result. The statement (2) follows similarly. □

Proposition 6.3. *If $(\mu, m) \in \hat{A}_k$, then*

$$Q_{k+\frac{1}{2}}(u, \mu) = -\frac{(u - |\mu| + 1 + \frac{z}{2})}{(u - |\mu| + \frac{z}{2})} \frac{\prod_{a \in A(\mu)} (u + c(a) - \frac{z}{2})}{\prod_{b \in R(\mu)} (u + c(b) - \frac{z}{2})} \tag{6.18}$$

and,

$$Q_{k+1}(u, \mu) = \frac{(u + |\mu| + 1 - \frac{z}{2})}{(u + |\mu| - \frac{z}{2})} \frac{\prod_{b \in R(\mu)} (u - c(b) + \frac{z}{2})}{\prod_{a \in A(\mu)} (u - c(a) + \frac{z}{2})}. \tag{6.19}$$

Proof. We prove (6.18). Let $(\mu, m) \in \hat{A}_k$. Since $f_{\mathfrak{t}^\mu}^{(k)}$ is a common eigenvector for the action of $\{Q_{i+\frac{1}{2}}(u) \mid i = 1, \dots, k\}$, the recursion (6.4) implies that

$$f_{\mathfrak{t}^\mu}^{(k)} Q_{i+\frac{1}{2}}(u) = Q_{\frac{1}{2}}(u) f_{\mathfrak{t}^\mu}^{(k)}, \qquad \text{for } i = 1, \dots, m.$$

Thus it is sufficient to consider the case where $\mu \vdash k$. Let $\mathfrak{t}^\mu = (\mu^{(0)}, \mu^{(\frac{1}{2})}, \dots, \mu^{(k)})$. We now verify that

$$f_{\mathfrak{t}^\mu}^{(k)} Q_{i+\frac{1}{2}}(u) = Q_{i+\frac{1}{2}}\left(u, \mu^{(i)}\right) f_{\mathfrak{t}^\mu}^{(k)} \qquad \text{for } i = 0, 1, \dots, k. \tag{6.20}$$

Let $i = 1, 2, \dots,$ and $\mu^{(i-1)} = \nu$ and $\mu^{(i)} = \lambda = \nu \cup \{\alpha\}$. Then $Q_{i+\frac{1}{2}}(u)$ acts on $f_{\mathfrak{t}^\mu}^{(k)}$ by the scalar

$$\begin{aligned} Q_{i-\frac{1}{2}}(u, \nu) \frac{(u + c_{\mathfrak{t}}(i))^2 - 1}{(u + c_{\mathfrak{t}}(i - \frac{1}{2}))^2 - 1} \frac{(u - c_{\mathfrak{t}}(i - \frac{1}{2}))^2}{(u + c_{\mathfrak{t}}(i))^2} &= -\frac{(u - |\nu| + 1 + \frac{z}{2})}{(u - |\nu| + \frac{z}{2})} \\ \times \frac{\prod_{a \in A(\nu)}(u + c(a) - \frac{z}{2})}{\prod_{b \in R(\nu)}(u + c(b) - \frac{z}{2})} \cdot \frac{(u + 1 + c(\alpha) + \frac{z}{2})(u - 1 + c(\alpha) - \frac{z}{2})}{(u - |\nu| - 1 + \frac{z}{2})(u - |\nu| + 1 + \frac{z}{2})} \frac{(u - |\nu| + \frac{z}{2})^2}{(u + c(\alpha) - \frac{z}{2})^2} \\ = -\frac{(u - |\lambda| + 1 + \frac{z}{2})}{(u - |\lambda| + \frac{z}{2})} \frac{\prod_{a \in A(\nu), a \neq \alpha}(u + c(a) - \frac{z}{2})}{\prod_{b \in R(\nu)}(u + c(b) - \frac{z}{2})} \frac{(u + 1 + c(\alpha) - \frac{z}{2})(u - 1 + c(\alpha) - \frac{z}{2})}{(u + c(\alpha) - \frac{z}{2})}. \end{aligned}$$

Let a_1, a_4 respectively be the nodes above and below α and a_2, a_3 respectively be the nodes to the right and left of α. Then $a_1 \in R(\nu) \Leftrightarrow a_2 \notin A(\lambda)$, and $a_3 \in R(\nu) \Leftrightarrow a_4 \notin A(\lambda)$. Since $c(a_1) = c(\alpha) + 1 = c(a_2)$ and $c(a_3) = c(\alpha) - 1 = c(a_4)$ the proof of (6.18) is complete. The statement (6.19) is verified similarly. ☐

7. Tables of Representing Matrices

The tables below give the representing matrices for the generators $p_i, p_{i+\frac{1}{2}}, \sigma_{i+\frac{1}{2}}, \sigma_{i+1}$ in selected representations of A_k and $A_{k+\frac{1}{2}}$ of small rank.

$A_1(z)$	$\lambda = \emptyset$
$p_1 \mapsto [z]$	

$A_{1+\frac{1}{2}}(z)$	$\lambda = (1)$
$p_1 \mapsto [0]$	
$p_{1+\frac{1}{2}} \mapsto [0]$	$\sigma_{1+\frac{1}{2}} \mapsto [1]$

$A_2(z)$	$\lambda = (1)$
$p_1 \mapsto \begin{bmatrix} z & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$	
$p_{1+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & \frac{z-1}{z^2} & 0 \\ 1 & \frac{z-1}{z} & 0 \\ 0 & 0 & 0 \end{bmatrix}$	$\sigma_{1+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$
$p_2 \mapsto \begin{bmatrix} 0 & 0 & 0 \\ 0 & \frac{z}{z-1} & \frac{z^2(z-2)}{(z-1)^2} \\ 0 & 1 & \frac{z(z-2)}{z-1} \end{bmatrix}$	$\sigma_2 \mapsto \begin{bmatrix} 0 & \frac{1}{z} & \frac{z-2}{z-1} \\ \frac{z}{z-1} & \frac{z-2}{z-1} & \frac{z(2-z)}{(z-1)^2} \\ 1 & -\frac{1}{z} & \frac{1}{z-1} \end{bmatrix}$

$A_{1+\frac{1}{2}}(z)$	$\lambda = \emptyset$
$p_1 \mapsto \begin{bmatrix} z & 0 \\ 0 & 0 \end{bmatrix}$	
$p_{1+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & \frac{z-1}{z^2} \\ 1 & \frac{z-1}{z} \end{bmatrix}$	$\sigma_{1+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$A_2(z)$	$\lambda = \emptyset$
$p_1 \mapsto \begin{bmatrix} z & 0 \\ 0 & 0 \end{bmatrix}$	
$p_{1+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & \frac{z-1}{z^2} \\ 1 & \frac{z-1}{z} \end{bmatrix}$	$\sigma_{1+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
$p_2 \mapsto \begin{bmatrix} z & 0 \\ 0 & 0 \end{bmatrix}$	$\sigma_2 \mapsto \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$A_2(z)$	$\lambda = (2)$
$p_1 \mapsto [0]$	
$p_{1+\frac{1}{2}} \mapsto [0]$	$\sigma_{1+\frac{1}{2}} \mapsto [1]$
$p_2 \mapsto [0]$	$\hat{\sigma}_2 \mapsto [1]$

$A_2(z)$	$\lambda = (1,1)$
$p_1 \mapsto [0]$	
$p_{1+\frac{1}{2}} \mapsto [0]$	$\sigma_{1+\frac{1}{2}} \mapsto [1]$
$p_2 \mapsto [0]$	$\sigma_2 \mapsto [-1]$

$A_{2+\frac{1}{2}}(z)$	$\lambda = \emptyset$
$p_1 \mapsto \begin{bmatrix} z & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & z & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	
$p_{1+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & \frac{z-1}{z^2} & 0 & 0 & 0 \\ 1 & \frac{z-1}{z} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{z} & \frac{z-1}{z^2} & 0 \\ 0 & 0 & 1 & \frac{z-1}{z} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	$\sigma_{1+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$p_2 \mapsto \begin{bmatrix} z & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{z}{z-1} & \frac{z^2(z-2)}{(z-1)^2} \\ 0 & 0 & 0 & 1 & \frac{z(z-2)}{z-1} \end{bmatrix}$	$\sigma_2 \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{z} & \frac{z-2}{z-1} \\ 0 & 0 & \frac{z}{z-1} & \frac{z-2}{z-1} & \frac{z(2-z)}{(z-1)^2} \\ 0 & 0 & 1 & -\frac{1}{z} & \frac{1}{z-1} \end{bmatrix}$
$p_{2+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & 0 & \frac{z-1}{z^2} & 0 & 0 \\ 0 & \frac{1}{z} & 0 & \frac{z-1}{z^2} & 0 \\ 1 & 0 & \frac{z-1}{z} & 0 & 0 \\ 0 & 1 & 0 & \frac{z-1}{z} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	$\sigma_{2+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{z} & \frac{z-2}{z-1} \\ 0 & 0 & 1 & 0 & 0 \\ 0 & \frac{z}{z-1} & 0 & \frac{z-2}{z-1} & \frac{z(2-z)}{(z-1)^2} \\ 0 & 1 & 0 & -\frac{1}{z} & \frac{1}{z-1} \end{bmatrix}$

$A_{2+\frac{1}{2}}(z)$	$\lambda=(1)$
$p_1 \mapsto \begin{bmatrix} z & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	
$p_{1+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & \frac{z-1}{z^2} & 0 & 0 & 0 \\ 1 & \frac{z-1}{z} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	$\sigma_{1+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$p_2 \mapsto \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{z}{z-1} & \frac{z^2(z-2)}{(z-1)^2} & 0 & 0 \\ 0 & 1 & \frac{z(z-2)}{z-1} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	$\sigma_2 \mapsto \begin{bmatrix} 0 & \frac{1}{z} & \frac{z-2}{z-1} & 0 & 0 \\ \frac{z}{z-1} & \frac{z-2}{z-1} & \frac{z(2-z)}{(z-1)^2} & 0 & 0 \\ 1 & -\frac{1}{z} & \frac{1}{z-1} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 \end{bmatrix}$
$p_{2+\frac{1}{2}} \mapsto \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{z-1}{z(z-2)} & \frac{(z-3)(z-1)}{z(z-2)^2} & \frac{(z-1)^2}{2z^2(z-2)} \\ 0 & 0 & \frac{1}{2} & \frac{z-3}{2(z-2)} & \frac{z-1}{4z} \\ 0 & 0 & 1 & \frac{z-3}{z-2} & \frac{z-1}{2z} \end{bmatrix}$	$\sigma_{2+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{z-1} & \frac{z}{(z-1)^2} & \frac{z(z-3)}{(z-1)(z-2)} & -\frac{1}{2} \\ 0 & \frac{z}{z(z-2)} & \frac{z^3-3z^2+2z-1}{z(z-2)(z-1)} & -\frac{z-3}{z(z-2)^2} & \frac{z-1}{2z^2(z-2)} \\ 0 & \frac{1}{2} & -\frac{1}{2(z-1)} & \frac{z-1}{2(z-2)} & \frac{z-1}{4z} \\ 0 & -1 & \frac{1}{z-1} & \frac{z-3}{z-2} & \frac{z+1}{2z} \end{bmatrix}$

$A_{2+\frac{1}{2}}(z)$	$\lambda = (2)$
$p_1 \mapsto [0]$	
$p_{1+\frac{1}{2}} \mapsto [0]$	$\sigma_{1+\frac{1}{2}} \mapsto [1]$
$p_2 \mapsto [0]$	$\sigma_2 \mapsto [1]$
$p_{+\frac{1}{2}} \mapsto [0]$	$\sigma_{2+\frac{1}{2}} \mapsto [1]$

$A_{2+\frac{1}{2}}(z)$	$\lambda = (1,1)$
$p_1 \mapsto [0]$	
$p_{1+\frac{1}{2}} \mapsto [0]$	$\sigma_{1+\frac{1}{2}} \mapsto [1]$
$p_2 \mapsto [0]$	$\sigma_2 \mapsto [-1]$
$p_{+\frac{1}{2}} \mapsto [0]$	$\sigma_{2+\frac{1}{2}} \mapsto [1]$

$A_3(z)$	$\lambda = \emptyset$
$p_1 \mapsto \begin{bmatrix} z & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & z & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	
$p_{1+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & \frac{z-1}{z^2} & 0 & 0 & 0 \\ 1 & \frac{z-1}{z} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{z} & \frac{z-1}{z^2} & 0 \\ 0 & 0 & 1 & \frac{z-1}{z} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	$\sigma_{1+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$
$p_2 \mapsto \begin{bmatrix} z & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{z}{z-1} & \frac{z^2(z-2)}{(z-1)^2} \\ 0 & 0 & 0 & 1 & \frac{z(z-2)}{z-1} \end{bmatrix}$	$\sigma_2 \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{z} & \frac{z-2}{z-1} \\ 0 & 0 & \frac{z}{z-1} & \frac{z-2}{z-1} & \frac{z(2-z)}{(z-1)^2} \\ 0 & 0 & 1 & -\frac{1}{z} & \frac{1}{z-1} \end{bmatrix}$
$p_{2+\frac{1}{2}} \mapsto \begin{bmatrix} \frac{1}{z} & 0 & \frac{z-1}{z^2} & 0 & 0 \\ 0 & \frac{1}{z} & 0 & \frac{z-1}{z^2} & 0 \\ 1 & 0 & \frac{z-1}{z} & 0 & 0 \\ 0 & 1 & 0 & \frac{z-1}{z} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	$\sigma_{2+\frac{1}{2}} \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{z} & \frac{z-2}{z-1} \\ 0 & 0 & 1 & 0 & 0 \\ 0 & \frac{z}{z-1} & 0 & \frac{z-2}{z-1} & \frac{z(2-z)}{(z-1)^2} \\ 0 & 1 & 0 & -\frac{1}{z} & \frac{1}{z-1} \end{bmatrix}$
$p_3 \mapsto \begin{bmatrix} z & 0 & 0 & 0 & 0 \\ 0 & z & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$	$\sigma_3 \mapsto \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$

References

[BB] A. Beliakova and C. Blanchet *Skein construction of idempotents in Birman–Murakami–Wenzl algebras*, Math. Ann. **321** (2001), 347–373.

[CO] J. Comes, and V. Ostrik, *On blocks of Deligne's category* $\underline{\mathrm{Rep}}(S_t)$, Adv. Math. **226** (2011), 1331–1377. MR2737787

[DW] W. Doran and D. Wales *The partition algebras revisited*, J. Algebra **231** (2000) 265–330. MR1779601 (2001i:16032)

[Ea] J. East, *Generators and relations for partition monoids and algebras*, preprint (2007).

[En] J. Enyang, *Jucys–Murphy elements and a presentation for partition algebras*, preprint (2010). arXiv:1009.1939v3

[En1] J. Enyang, *Cellular bases for algebras with a Jones basic construction*, preprint (2011). arXiv:1106.5339

[En2] J. Enyang, *Gram determinants for algebras with a Jones basic construction*, preprint (2011).

[GG] F.M. Goodman and J. Graber, *Cellularity and the Jones basic construction*, Adv. in App. Math. **46** (2011) 312–362. doi:10.1016/j.aam.2010.10.003

[GG1] F.M. Goodman and J. Graber, *On cellular algebras with Jucys–Murphy elements*, J. Algebra **330**, 147–176 (2011). doi:10.1016/j.jalgebra.2011.01.002

[GL] J. Graham and G. Lehrer, *Cellular algebras*, Invent. Math., **123** (1996), 1–34. MR1376244 (97h:20016)

[HR] T. Halverson and A. Ram, *Partition algebras*, European J. Combin. **26** (2005), 869–921. MR2552957 (2010m:16047)

[Jo] V.F.R Jones, *The Potts model and the symmetric group,* in: Subfactors: Proceedings of the Taniguchi Symposium on Operator Algebras (Kuzeso, 1993), River Edge, NJ, World Sci. Publishing, 1994. MR1317365 (97b:82023)

[Kl] A. Kleshchev, *Linear and projective representations of symmetric groups,* Cambridge Tracts in Mathematics, 163. Cambridge University Press, Cambridge, 2005. MR2165457 (2007b:20022)

[Ko] M. Kosuda, *Irreducible representations of the party algebra,* Osaka J. Math. **43** (2006), 431–474. MR2262344 (2007g:05190)

[Ko1] M. Kosuda, *The partition algebra, its characterization and representations,* Ryukyu Math. J. **22** (2009), 1–29. MR2583260 (2011c:16083)

[LR] R. Leduc and A. Ram, *A ribbon Hopf algebra approach to the irreducible representations of centralizer algebras: the Brauer, Birman-Wenzl, and type A Iwahori-Hecke algebras*, Adv. Math. **125** (1997), 1–94. MR1427801 (98c:20015)

[Mar] P. Martin, *Potts models and related problems in statistical mechanics,* Series on Advances in Statistical Mechanics, vol. 5, World Scientific Publishing Co. Inc., Teaneck NJ, 1991. MR1103994 (92m:82030)

[Mar1] P. Martin, *Temperley-Lieb algebras for nonplanar statistical mechanics – the partition algebra construction,* J. Knot Theory Ramifications 3 (1994), 51–82. MR1265453 (95a:82022)

[Mar2] P. Martin, *The partition algebra and the Potts model transfer matrix spectrum in high dimensions,* J. Phys. A 33 (2000), 3669–3695.

[Mat] A. Mathas, *Seminormal forms and Gram determinants for cellular algebras*, J. Reine Angew. Math., **619** (2008), 141–173. With an appendix by M. Soriano. MR2414949 (2009e:16059)

[Na] M. Nazarov, *Young's orthogonal form for Brauer's centralizer algebra*, J. Algebra, **182** (1996), 664–693. MR1398116 (97m:20057)

[VO] A. Vershik and A. Okounkov, *A new approach to representation theory of symmetric groups II,* J. Math. Sci. (N. Y.) **131** (2005), 5471–5494. MR2050688 (2005c:20024)

[Xi] C.C. Xi, *Partition algebras are cellular*, Comp. Math., **119** (1999), 99–109. MR1711582 (2000i:16022)

[Yo] A. Young, *On quantitative substitutional analysis, IV*, Proc. Lond. Math. Soc. **31** (1931), 253–289.

Department of Mathematics and Statistics, University of Melbourne, Parkville VIC 3010, Australia

E-mail address: `jenyang@unimelb.edu.au`